

- Aarsland, Dag; Sardahaee, Farzaneh S.; Anderssen, Sigmund; Ballard, Clive (2010): Is physical activity a potential preventive factor for vascular dementia? A systematic review. In: *Aging & Mental Health* 14 (4), S. 386–395. DOI: 10.1080/13607860903586136.
- Abbott, Robert D.; White, Lon R.; Ross, G. Webster; Masaki, Kamal H.; Curb, J. David; Petrovitch, Helen (2004): Walking and dementia in physically capable elderly men. In: *JAMA* 292 (12), S. 1447–1453. DOI: 10.1001/jama.292.12.1447.
- Abramson, Jerome L.; Weintraub, William S.; Vaccarino, Viola (2002): Association between pulse pressure and C-reactive protein among apparently healthy US adults. In: *Hypertension* 39 (2), S. 197–202. DOI: 10.1161/hy0202.104270.
- ACIL, A. A.; DOGAN, S.; DOGAN, O. (2008): The effects of physical exercises to mental state and quality of life in patients with schizophrenia. In: *Journal of Psychiatric and Mental Health Nursing* 15 (10), S. 808–815. DOI: 10.1111/j.1365-2850.2008.01317.x.
- Adamopoulos, Stamatis; Coats, Andrew J.S.; Brunotte, François; Arnolda, Leonard; Meyer, Theo; Thompson, Campbell H. et al. (1993): Physical training improves skeletal muscle metabolism in patients with chronic heart failure. In: *Journal of the American College of Cardiology* 21 (5), S. 1101–1106. DOI: 10.1016/0735-1097(93)90231-O.
- Adamopoulos, S.; Parissis, J.; Kroupis, C.; Georgiadis, M.; Karatzas, D.; Karavolias, G. et al. (2001): Physical training reduces peripheral markers of inflammation in patients with chronic heart failure. In: *European Heart Journal* 22 (9), S. 791–797. DOI: 10.1053/euhj.2000.2285.
- Adamopoulos, Stamatis; Parissis, John; Karatzas, Dimitrios; Kroupis, Christos; Georgiadis, Michael; Karavolias, George et al. (2002): Physical training modulates proinflammatory cytokines and the soluble Fas/soluble Fasligand system in patients with chronic heart failure. In: *Journal of the American College of Cardiology* 39 (4), S. 653–663. DOI: 10.1016/S0735-1097(01)01795-8.
- Adamsen, Lis; Quist, Morten; Andersen, Christina; Møller, Tom; Herrstedt, Jørn; Kronborg, Dorte et al. (2009): Effect of a multimodal high intensity exercise intervention in cancer patients undergoing chemotherapy: randomised controlled trial. In: *BMJ* 339 (oct13 1), b3410. DOI: 10.1136/bmj.b3410.
- Ades, P. A. (2001): Cardiac rehabilitation and secondary prevention of coronary heart disease. In: *The New England journal of medicine* 345 (12), S. 892–902. DOI: 10.1056/NEJMra001529.
- Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies (2002). In: *The Lancet* 360 (9349), S. 1903–1913.
- Ahlskog, J. Eric (2011): Does vigorous exercise have a neuroprotective effect in Parkinson disease? In: *Neurology* 77 (3), S. 288–294. DOI: 10.1212/WNL.0b013e318225ab66.
- Akbaraly, T. N.; Portet, F.; Fustinoni, S.; Dartigues, J-F; Artero, S.; Rouaud, O. et al. (2009): Leisure activities and the risk of dementia in the elderly: results from the Three-City Study. In: *Neurology* 73 (11), S. 854–861. DOI: 10.1212/WNL.0b013e3181b7849b.
- Alam, Saima; Stolinski, Michael; Pentecost, Claire; Boroujerdi, Massoud A.; Jones, Richard H.; Sonksen, Peter H.; Umpleby, A. Margot (2004): The effect of a six-month exercise program on very low-density lipoprotein apolipoprotein B secretion in type 2 diabetes. In: *The Journal of clinical endocrinology and metabolism* 89 (2), S. 688–694. DOI: 10.1210/jc.2003-031036.
- Alamanos, Yannis; Voulgari, Paraskevi V.; Drosos, Alexandros A. (2006): Incidence and prevalence of rheumatoid arthritis, based on the 1987 American College of Rheumatology criteria: a systematic review. In: *Seminars in Arthritis and Rheumatism* 36 (3), S. 182–188. DOI: 10.1016/j.semarthrit.2006.08.006.
- Albright, A.; Franz, M.; Hornsby, G.; Kriska, A.; Marrero, D.; Ullrich, I.; Verity, L. S. (2000): American College of Sports Medicine position stand. Exercise and type 2 diabetes. In: *Medicine & Science in Sports & Exercise* 32 (7), S. 1345–1360. DOI: 10.1097/00005768-200007000-00024.
- Alfano, Catherine M.; Smith, Ashley Wilder; Irwin, Melinda L.; Bowen, Deborah J.; Sorensen, Bess; Reeve, Bryce B. et al. (2007): Physical activity, long-term symptoms, and physical health-related quality of life among breast cancer survivors: a prospective analysis. In: *Journal of Cancer Survivorship* 1 (2), S. 116–128. DOI: 10.1007/s11764-007-0014-1.
- Allenberg, K.; Johansen, K.; Saltin, B. (1988): Skeletal muscle adaptations to physical training in type II (non-insulin-dependent) diabetes mellitus. In: *Acta Medica Scandinavica* 223 (4), S. 365–373. DOI: 10.1111/j.0954-6820.1988.tb15886.x.
- Alonso-Frech, Fernando; Sanahuja, Juan Juni; Rodriguez, Amelia Mendoza (2011): Exercise and physical therapy in early management of Parkinson disease. In: *The Neurologist* 17 (6 Suppl 1), S47-53. DOI: 10.1097/NRL.0b013e31823968ec.
- Amatya (2013): Non pharmacological interventions for spasticity in multiple sclerosis. In: *Cochrane Database Syst Rev* 2, CD009974.
- American Diabetes Association (2002): Clinical practice recommendations. In: *Diabetes Care* Jan, S1.
- American Thoracic Society (1999): Pulmonary rehabilitation - 1999. Official statement of the American Thoracic Society, November 1998. In: *Am J Respir Crit Care Med* 1999 (159), S. 1666.
- Andel, Ross; Crowe, Michael; Pedersen, Nancy L.; Fratiglioni, Laura; Johansson, Boo; Gatz, Margaret (2008): Physical exercise at midlife and risk of dementia three decades later: a population-based study of Swedish twins. In: *The Journals of Gerontology: Series A* 63 (1), S. 62–66. DOI: 10.1093/gerona/63.1.62.

- Anderiesen, Hester; Scherder, Erik J. A.; Goossens, Richard H. M.; Sonneveld, Marieke H. (2014): A systematic review--physical activity in dementia: the influence of the nursing home environment. In: *Applied Ergonomics* 45 (6), S. 1678–1686. DOI: 10.1016/j.apergo.2014.05.011.
- Andersen, R. E.; Wadden, T. A.; Bartlett, S. J.; Zemel, B.; Verde, T. J.; Franckowiak, S. C. (1999): Effects of lifestyle activity vs structured aerobic exercise in obese women: a randomized trial. In: *JAMA* 281 (4), S. 335–340. DOI: 10.1001/jama.281.4.335.
- Anderson, J. W.; Konz, E. C.; Frederich, R. C.; Wood, C. L. (2001): Long-term weight-loss maintenance: a meta-analysis of US studies. In: *The American Journal of Clinical Nutrition* 74 (5), S. 579–584. DOI: 10.1093/ajcn/74.5.579.
- Anker, S. D.; Chua, T. P.; Ponikowski, P.; Harrington, D.; Swan, J. W.; Kox, W. J. et al. (1997): Hormonal changes and catabolic/anabolic imbalance in chronic heart failure and their importance for cardiac cachexia. In: *Circulation* 96 (2), S. 526–534. DOI: 10.1161/01.CIR.96.2.526.
- Armstrong, Neil; Simons-Morton, Bruce (1994): Physical Activity and Blood Lipids in Adolescents. In: *Pediatric Exercise Science* 6 (4), S. 381–405. DOI: 10.1123/pes.6.4.381.
- Arnold, Cathy M.; Faulkner, Robert A. (2010): The effect of aquatic exercise and education on lowering fall risk in older adults with hip osteoarthritis. In: *Journal of Aging and Physical Activity* 18 (3), S. 245–260. DOI: 10.1123/japa.18.3.245.
- Babiyak, M.; Blumenthal, J. A.; Herman, S.; Khatri, P.; Doraiswamy, M.; Moore, K. et al. (2000): Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. In: *Psychosomatic Medicine* 62 (5), S. 633–638. DOI: 10.1097/00006842-200009000-00006.
- Bahrke, Michael S.; Morgan, William P. (1978): Anxiety reduction following exercise and meditation. In: *Cogn Ther Res* 2 (4), S. 323–333. DOI: 10.1007/BF01172650.
- Baillet, Athan; Zeboulon, Nadine; Gossec, Laure; Combescurre, Christophe; Bodin, Louis-Antoine; Juvin, Robert et al. (2010): Efficacy of cardiorespiratory aerobic exercise in rheumatoid arthritis: meta-analysis of randomized controlled trials. In: *Arthritis Care & Research* 62 (7), S. 984–992. DOI: 10.1002/acr.20146.
- Baker, Laura D.; Frank, Laura L.; Foster-Schubert, Karen; Green, Pattie S.; Wilkinson, Charles W.; McTiernan, Anne et al. (2010): Effects of aerobic exercise on mild cognitive impairment: a controlled trial. In: *Archives of Neurology* 67 (1), S. 71–79. DOI: 10.1001/archneurol.2009.307.
- Baker, N. A.; Tickle-Degnen, L. (2001): The effectiveness of physical, psychological, and functional interventions in treating clients with multiple sclerosis: a meta-analysis. In: *American Journal of Occupational Therapy* 55 (3), S. 324–331. DOI: 10.5014/ajot.55.3.324.
- Barefoot, J. C.; Heitmann, B. L.; Helms, M. J.; Williams, R. B.; Surwit, R. S.; Siegler, I. C. (1998): Symptoms of depression and changes in body weight from adolescence to mid-life. In: *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 22 (7), S. 688–694. DOI: 10.1038/sj.jjo.0800647.
- Barnard, R. James; Ugianskis, Erika J.; Martin, Deborah A.; Inkeles, Stephen B. (1992): Role of diet and exercise in the management of hyperinsulinemia and associated atherosclerotic risk factors. In: *The American Journal of Cardiology* 69 (5), S. 440–444. DOI: 10.1016/0002-9149(92)90981-4.
- Barry, Vaughn W.; Baruth, Meghan; Beets, Michael W.; Durstine, J. Larry; Liu, Jihong; Blair, Steven N. (2014): Fitness vs. fatness on all-cause mortality: a meta-analysis. In: *Progress in Cardiovascular Diseases* 56 (4), S. 382–390. DOI: 10.1016/j.pcad.2013.09.002.
- Bartley, Christine A.; Hay, Madeleine; Bloch, Michael H. (2013): Meta-analysis: aerobic exercise for the treatment of anxiety disorders. In: *Progress in neuro-psychopharmacology & biological psychiatry* 45, S. 34–39. DOI: 10.1016/j.pnpbp.2013.04.016.
- Baslund, B.; Lyngberg, K.; Andersen, V.; Halkjaer Kristensen, J.; Hansen, M.; Klokke, M.; Pedersen, B. K. (1993): Effect of 8 wk of bicycle training on the immune system of patients with rheumatoid arthritis. In: *Journal of applied physiology (Bethesda, Md. : 1985)* 75 (4), S. 1691–1695. DOI: 10.1152/jappl.1993.75.4.1691.
- Battegay, Edouard; Lip, Gregory Y. H.; Bakris, George L. (2005): Hypertension. Principles and practice. Boca Raton: Taylor & Francis (Hypertension).
- Bauldoff, Gerene S.; Hoffman, Leslie A.; Zullo, Thomas G.; Sciarba, Frank C. (2002): Exercise maintenance following pulmonary rehabilitation: effect of distractive stimuli. In: *Chest* 122 (3), S. 948–954. DOI: 10.1378/chest.122.3.948.
- Bautch, J. C.; Clayton, M. K.; Chu, Q.; Johnson, K. A. (2000): Synovial fluid chondroitin sulphate epitopes 3B3 and 7D4, and glycosaminoglycan in human knee osteoarthritis after exercise. In: *Annals of the Rheumatic Diseases* 59 (11), S. 887–891. DOI: 10.1136/ard.59.11.887.
- Bautmans, Ivan; van Arken, Judith; van Mackelenberg, Mike; Mets, Tony (2010): Rehabilitation using manual mobilization for thoracic kyphosis in elderly postmenopausal patients with osteoporosis. In: *Journal of Rehabilitation Medicine* 42 (2), S. 129–135. DOI: 10.2340/16501977-0486.
- Beals (1985): Measurement of exercise tolerance in patients with rheumatoid arthritis and osteoarthritis. In: *J Rheumatol* 12, S. 458.

- Beck-Nielsen H Henriksen JE Hermansen K Madsen LD Olivarius NF Mandrup-Poulsen TR Pedersen OB Richelsen B Schmitz OE [Type 2 diabetes and the metabolic syndrome - diagnosis and treatment] Copenhagen Laegeforeningens forlag 2000 1 36 (2000).
- Becofsky, Katie M.; Sui, Xuemei; Lee, Duck-chul; Wilcox, Sara; Zhang, Jiajia; Blair, Steven N. (2015): A prospective study of fitness, fatness, and depressive symptoms. In: *American Journal of Epidemiology* 181 (5), S. 311–320. DOI: 10.1093/aje/kwu330.
- Beebe, Lora Humphrey; Tian, Lili; Morris, Nancy; Goodwin, Ann; Allen, Staccie Swant; Kuldau, John (2005): Effects of exercise on mental and physical health parameters of persons with schizophrenia. In: *Issues in Mental Health Nursing* 26 (6), S. 661–676. DOI: 10.1080/01612840590959551.
- Beggs (2013): Swimming training for asthma in children and adolescents aged 18 years and under. In: *Cochrane Database Syst Rev* 4, CD009607.
- Beling, Janna; Roller, Margaret (2009): Multifactorial intervention with balance training as a core component among fall-prone older adults. In: *Journal of Geriatric Physical Therapy* 32 (3), S. 125–133. DOI: 10.1519/00139143-200932030-00008.
- Benatti, Fabiana B.; Pedersen, Bente K. (2015): Exercise as an anti-inflammatory therapy for rheumatic diseases-myokine regulation. In: *Nature Reviews Rheumatology* 11 (2), S. 86–97. DOI: 10.1038/nrrheum.2014.193.
- Bendermacher (2006): Supervised exercise therapy versus non-supervised exercise therapy for intermittent claudication. In: *Cochrane Database Syst Rev* 19, CD005263.
- Bennell, K. L.; Hunt, M. A.; Wrigley, T. V.; Hunter, D. J.; McManus, F. J.; Hodges, P. W. et al. (2010): Hip strengthening reduces symptoms but not knee load in people with medial knee osteoarthritis and varus malalignment: a randomised controlled trial. In: *Osteoarthritis and Cartilage* 18 (5), S. 621–628. DOI: 10.1016/j.joca.2010.01.010.
- Beraki, Å.; Magnuson, A.; Särnblad, S.; Åman, J.; Samuelsson, U. (2014): Increase in physical activity is associated with lower HbA1c levels in children and adolescents with type 1 diabetes: results from a cross-sectional study based on the Swedish pediatric diabetes quality registry (SWEDIABKIDS). In: *Diabetes Research and Clinical Practice* 105 (1), S. 119–125. DOI: 10.1016/j.diabres.2014.01.029.
- Beyer, Nina; Simonsen, Lene; Bülow, Jens; Lorenzen, Tove; Jensen, Dorte V.; Larsen, Lone et al. (2007): Old women with a recent fall history show improved muscle strength and function sustained for six months after finishing training. In: *Aging Clinical and Experimental Research* 19 (4), S. 300–309. DOI: 10.1007/BF03324706.
- Bild (1996): Correlates and predictors of weight loss in young adults: the CARDIA study. In: *Int J Obes Relat Metab Disord* 20, S. 47.
- Bjørnebekk, Astrid; Mathé, Aleksander A.; Brené, Stefan (2005): The antidepressant effect of running is associated with increased hippocampal cell proliferation. In: *The International Journal of Neuropsychopharmacology* 8 (3), S. 357–368. DOI: 10.1017/S1461145705005122.
- Blood Pressure Lowering Treatment Trialists' Collaboration (2000): Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. In: *Lancet* 355, S. 1955.
- Blumenthal, J. A.; Babyak, M. A.; Moore, K. A.; Craighead, W. E.; Herman, S.; Khatri, P. et al. (1999): Effects of exercise training on older patients with major depression. In: *Archives of Internal Medicine* 159 (19), S. 2349–2356. DOI: 10.1001/archinte.159.19.2349.
- Bogardus, C.; Ravussin, E.; Robbins, D. C.; Wolfe, R. R.; Horton, E. S.; Sims, E. A. (1984): Effects of physical training and diet therapy on carbohydrate metabolism in patients with glucose intolerance and non-insulin-dependent diabetes mellitus. In: *Diabetes* 33 (4), S. 311–318. DOI: 10.2337/diab.33.4.311.
- Booth (2012): Lack of exercise is a major cause of chronic diseases. In: *Compr Physiol* 2, S. 1143.
- Borer, Katarina T. (2005): Physical activity in the prevention and amelioration of osteoporosis in women : interaction of mechanical, hormonal and dietary factors. In: *Sports Medicine* 35 (9), S. 779–830. DOI: 10.2165/00007256-200535090-00004.
- Bosscher (1993): Running and mixed physical exercises with depressed psychiatric patients. In: *Int J Sport Psychol* 24, S. 170.
- Boudou, P.; Sobngwi, E.; Mauvais-Jarvis, F.; Vexiau, P.; Gautier, J-F (2003): Absence of exercise-induced variations in adiponectin levels despite decreased abdominal adiposity and improved insulin sensitivity in type 2 diabetic men. In: *European Journal of Endocrinology* 149 (5), S. 421–424. DOI: 10.1530/eje.0.1490421.
- Boulé, N. G.; Haddad, E.; Kenny, G. P.; Wells, G. A.; Sigal, R. J. (2001): Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: a meta-analysis of controlled clinical trials. In: *JAMA* 286 (10), S. 1218–1227. DOI: 10.1001/jama.286.10.1218.
- Boulé, N. G.; Kenny, G. P.; Haddad, E.; Wells, G. A.; Sigal, R. J. (2003): Meta-analysis of the effect of structured exercise training on cardiorespiratory fitness in Type 2 diabetes mellitus. In: *Diabetologia* 46 (8), S. 1071–1081. DOI: 10.1007/s00125-003-1160-2.
- Boyko, E. J.; Fujimoto, W. Y.; Leonetti, D. L.; Newell-Morris, L. (2000): Visceral adiposity and risk of type 2 diabetes: a prospective study among Japanese Americans. In: *Diabetes Care* 23 (4), S. 465–471. DOI: 10.2337/diacare.23.4.465.
- Boysen, Gudrun; Krarup, Lars-Henrik (2009): Benefits of physical activity for stroke survivors. In: *Expert Review of Neurotherapeutics* 9 (2), S. 147–149. DOI: 10.1586/14737175.9.2.147.

- Bradham, William S.; Moe, Gordon; Wendt, Katherine A.; Scott, Amelia A.; Konig, Andrea; Romanova, Marina et al. (2002): TNF-alpha and myocardial matrix metalloproteinases in heart failure: relationship to LV remodeling. In: *American journal of physiology. Heart and circulatory physiology* 282 (4), H1288-95. DOI: 10.1152/ajpheart.00526.2001.
- Bradley (2002): Physical training for cystic fibrosis. In: *Cochrane Database Syst Rev*, S. 1.
- Brandon, L. Jerome; Gaasch, Debbi A.; Boyette, Lisa W.; Lloyd, Adrienne M. (2003): Effects of long-term resistive training on mobility and strength in older adults with diabetes. In: *The Journals of Gerontology: Series A* 58 (8), S. 740-745. DOI: 10.1093/gerona/58.8.M740.
- Braunwald (2008): Braunwald's heart disease: a textbook of cardiovascular medicine.
- BRENNAN, F. M.; MAINI, R. N.; FELDMANN, M. (1992): TNF alpha--a pivotal role in rheumatoid arthritis? In: *British journal of rheumatology* 31 (5), S. 293-298. DOI: 10.1093/rheumatology/31.5.293.
- Brinkmann, J. R.; Hoskins, T. A. (1979): Physical conditioning and altered self-concept in rehabilitated hemiplegic patients. In: *Physical Therapy* 59 (7), S. 859-865. DOI: 10.1093/ptj/59.7.859.
- Briscoe, Vanessa Jones; Tate, Donna Bowman; Davis, Stephen Neil (2007): Type 1 diabetes: exercise and hypoglycemia. In: *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme* 32 (3), S. 576-582. DOI: 10.1139/H07-025.
- British Thoracic Society Standards of Care Subcommittee on Pulmonary Rehabilitation (2001): Pulmonary rehabilitation. In: *Thorax* 56 (11), S. 827-834. DOI: 10.1136/thorax.56.11.827.
- Broderick, J. P.; Adams, H. P.; Barsan, W.; Feinberg, W.; Feldmann, E.; Grotta, J. et al. (1999): Guidelines for the management of spontaneous intracerebral hemorrhage: A statement for healthcare professionals from a special writing group of the Stroke Council, American Heart Association. In: *Stroke* 30 (4), S. 905-915. DOI: 10.1161/01.STR.30.4.905.
- Brooks (1995): Exercise physiology: human bioenergetics and its applicatons.
- BROWN, S.; BIRTWISTLE, J.; ROE, L.; THOMPSON, C. (1999): The unhealthy lifestyle of people with schizophrenia. In: *Psychological Medicine* 29 (3), S. 697-701. DOI: 10.1017/S0033291798008186.
- Brown, David R.; Pate, Russell R.; Pratt, Michael; Wheeler, Fran; Buchner, David; Ainsworth, Barbara; Macera, Carol (2001): Physical Activity and Public Health: Training courses for researchers and practitioners. In: *Public Health Reports* 116 (3), S. 197-202. DOI: 10.1016/S0033-3549(04)50034-1.
- Brox, Jens Ivar; Reikerås, Olav; Nygaard, Øystein; Sørensen, Roger; Indahl, Aage; Holm, Inger et al. (2006): Lumbar instrumented fusion compared with cognitive intervention and exercises in patients with chronic back pain after previous surgery for disc herniation: a prospective randomized controlled study. In: *Pain* 122 (1-2), S. 145-155. DOI: 10.1016/j.pain.2006.01.027.
- Brox, Jens Ivar; Sørensen, Roger; Friis, Astrid; Nygaard, Øystein; Indahl, Aage; Keller, Anne et al. (2003): Randomized clinical trial of lumbar instrumented fusion and cognitive intervention and exercises in patients with chronic low back pain and disc degeneration. In: *Spine* 28 (17), S. 1913-1921. DOI: 10.1097/01.BRS.0000083234.62751.7A.
- Budson, Andrew E.; Solomon, Paul R. (2011): Memory Loss: A Practical Guide for Clinicians. 1. Aufl.: Elsevier Health Sciences (Memory Loss).
- Burke, Thomaz Nogueira; França, Fábio Jorge Renovato; Ferreira de Meneses, Sarah Rúbica; Cardoso, Viviam Inhasz; Marques, Amélia Pasqual (2010): Postural control in elderly persons with osteoporosis: Efficacy of an intervention program to improve balance and muscle strength: a randomized controlled trial. In: *American Journal of Physical Medicine & Rehabilitation* 89 (7), S. 549-556. DOI: 10.1097/PHM.0b013e3181ddcccd2.
- Burt, V. L.; Whelton, P.; Roccella, E. J.; Brown, C.; Cutler, J. A.; Higgins, M. et al. (1995): Prevalence of hypertension in the US adult population. Results from the Third National Health and Nutrition Examination Survey, 1988-1991. In: *Hypertension* 25 (3), S. 305-313. DOI: 10.1161/01.HYP.25.3.305.
- Cakar (2010): Jumping combined exercise programs reduce fall risk and improve balance and life quality of elderly people who live in a long-term care facility. In: *Eur J Phys Rehabil Med* 46, S. 59.
- Calver, A.; Collier, J.; Vallance, P. (1992): Inhibition and stimulation of nitric oxide synthesis in the human forearm arterial bed of patients with insulin-dependent diabetes. In: *Journal of Clinical Investigation* 90 (6), S. 2548-2554. DOI: 10.1172/JCI116149.
- Campbell, R. Keith (2009): Type 2 diabetes: where we are today: an overview of disease burden, current treatments, and treatment strategies. In: *Journal of the American Pharmacists Association : JAPhA* 49 Suppl 1 (5), S3-9. DOI: 10.1331/JAPhA.2009.09077.
- Campbell Burton, C. Alexia; Murray, Jenni; Holmes, John; Astin, Felicity; Greenwood, Darren; Knapp, Peter (2013): Frequency of anxiety after stroke: a systematic review and meta-analysis of observational studies. In: *International Journal of Stroke* 8 (7), S. 545-559. DOI: 10.1111/j.1747-4949.2012.00906.x.
- Campbell, Stephanie; Marriott, Michael; Nahmias, Claude; MacQueen, Glenda M. (2004): Lower hippocampal volume in patients suffering from depression: a meta-analysis. In: *The American journal of psychiatry* 161 (4), S. 598-607. DOI: 10.1176/appi.ajp.161.4.598.

- Carda, Stefano; Cisari, Carlo; Invernizzi, Marco; Bevilacqua, Maurizio (2009): Osteoporosis after stroke: a review of the causes and potential treatments. In: *Cerebrovascular diseases (Basel, Switzerland)* 28 (2), S. 191–200. DOI: 10.1159/000226578.
- Carlsen, Kai-Håkon; Carlsen, Karin C.L. (2002): Exercise-induced asthma. In: *Paediatric Respiratory Reviews* 3 (2), S. 154–160. DOI: 10.1016/S1526-0550(02)00009-4.
- Carlson, Debra J.; Dieberg, Gudrun; Hess, Nicole C.; Millar, Philip J.; Smart, Neil A. (2014): Isometric exercise training for blood pressure management: a systematic review and meta-analysis. In: *Mayo Clinic Proceedings* 89 (3), S. 327–334. DOI: 10.1016/j.mayocp.2013.10.030.
- Carral, Florentino; Gutiérrez, José Vicente; Ayala, María del Carmen; García, Gloria; Aguilar, Manuel (2013): Intense physical activity is associated with better metabolic control in patients with type 1 diabetes. In: *Diabetes Research and Clinical Practice* 101 (1), S. 45–49. DOI: 10.1016/j.diabres.2013.04.006.
- Carson (2013): Physical training for asthma. In: *Cochrane Database Syst Rev* 9, CD001116.
- Carter (2002): Community-based exercise program reduces risk factors for falls in 65- to 75-year-old women with osteoporosis: randomized controlled trial. In: *CMAJ* 167, S. 997.
- Carter, N. D.; Khan, K. M.; Petit, M. A.; Heinonen, A.; Waterman, C.; Donaldson, M. G. et al. (2001): Results of a 10 week community based strength and balance training programme to reduce fall risk factors: a randomised controlled trial in 65-75 year old women with osteoporosis. In: *British Journal of Sports Medicine* 35 (5), S. 348–351. DOI: 10.1136/bjism.35.5.348.
- Chaddock, Laura; Erickson, Kirk I.; Prakash, Ruchika Shaurya; Kim, Jennifer S.; Voss, Michelle W.; VanPatter, Matt et al. (2010): A neuroimaging investigation of the association between aerobic fitness, hippocampal volume, and memory performance in preadolescent children. In: *Brain Research* 1358, S. 172–183. DOI: 10.1016/j.brainres.2010.08.049.
- Chamove, A. S. (1986): Positive short-term effects of activity on behaviour in chronic schizophrenic patients. In: *British Journal of Clinical Psychology* 25 (Pt 2) (2), S. 125–133. DOI: 10.1111/j.2044-8260.1986.tb00681.x.
- Chen, Xiaoli; Zheng, Ying; Zheng, Wei; Gu, Kai; Chen, Zhi; Lu, Wei; Shu, Xiao Ou (2009): The effect of regular exercise on quality of life among breast cancer survivors. In: *American Journal of Epidemiology* 170 (7), S. 854–862. DOI: 10.1093/aje/kwp209.
- Cho, Eo Rin; Shin, Aesun; Kim, Jeongseon; Jee, Sun Ha; Sung, Joohon (2009): Leisure-time physical activity is associated with a reduced risk for metabolic syndrome. In: *Annals of Epidemiology* 19 (11), S. 784–792. DOI: 10.1016/j.annepidem.2009.06.010.
- Chong, P. F.; Gollidge, J.; Greenhalgh, R. M.; Davies, A. H. (2000): Exercise therapy or angioplasty? A summation analysis. In: *European Journal of Vascular and Endovascular Surgery* 20 (1), S. 4–12. DOI: 10.1053/ejvs.2000.1112.
- Church, Timothy S.; Blair, Steven N.; Cocreham, Shannon; Johannsen, Neil; Johnson, William; Kramer, Kimberly et al. (2010): Effects of aerobic and resistance training on hemoglobin A1c levels in patients with type 2 diabetes: a randomized controlled trial. In: *JAMA* 304 (20), S. 2253–2262. DOI: 10.1001/jama.2010.1710.
- Clark, C. J.; Cochrane, L. M. (1988): Assessment of work performance in asthma for determination of cardiorespiratory fitness and training capacity. In: *Thorax* 43 (10), S. 745–749. DOI: 10.1136/thx.43.10.745.
- Clarkson, Peter; Celermajer, David S.; Donald, Ann E.; Sampson, Mike; Sorensen, Keld E.; Adams, Mark et al. (1996): Impaired vascular reactivity in insulin-dependent diabetes mellitus is related to disease duration and low density lipoprotein cholesterol levels. In: *Journal of the American College of Cardiology* 28 (3), S. 573–579. DOI: 10.1016/0735-1097(96)82380-1.
- Coakley, E. H.; Rimm, E. B.; Colditz, G.; Kawachi, I.; Willett, W. (1998): Predictors of weight change in men: results from the Health Professionals Follow-up Study. In: *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 22 (2), S. 89–96. DOI: 10.1038/sj.ijo.0800549.
- COATS, A. (1990): Effects of physical training in chronic heart failure. In: *The Lancet* 335 (8681), S. 63–66. DOI: 10.1016/0140-6736(90)90536-E.
- Coats, A. J.; Adamopoulos, S.; Radaelli, A.; McCance, A.; Meyer, T. E.; Bernardi, L. et al. (1992): Controlled trial of physical training in chronic heart failure. Exercise performance, hemodynamics, ventilation, and autonomic function. In: *Circulation* 85 (6), S. 2119–2131. DOI: 10.1161/01.CIR.85.6.2119.
- Coggan (1993): Effect of prolonged exercise on muscle citrate concentration before and after endurance training in men. In: *Am J Physiol* 264, E215.
- Collins, R.; MacMahon, S. (1994): Blood pressure, antihypertensive drug treatment and the risks of stroke and of coronary heart disease. In: *British Medical Bulletin* 50 (2), S. 272–298. DOI: 10.1093/oxfordjournals.bmb.a072892.
- Collins, R.; Peto, R.; MacMahon, S.; Godwin, J.; Qizilbash, N.; Hebert, P. et al. (1990): Blood pressure, stroke, and coronary heart disease. In: *The Lancet* 335 (8693), S. 827–838. DOI: 10.1016/0140-6736(90)90944-Z.
- Combs, Stephanie A.; Diehl, M. Dyer; Chrzastowski, Casey; Didrick, Nora; McCoin, Brittany; Mox, Nicholas et al. (2013): Community-based group exercise for persons with Parkinson disease: a randomized controlled trial. In: *NeuroRehabilitation* 32 (1), S. 117–124. DOI: 10.3233/NRE-130828.
- Conn, Vicki S. (2010): Anxiety outcomes after physical activity interventions: meta-analysis findings. In: *Nursing Research* 59 (3), S. 224–231. DOI: 10.1097/NNR.0b013e3181dbb2f8.

- Conn, V. S.; Hafdahl, A. R.; Mehr, D. R.; LeMaster, J. W.; Brown, S. A.; Nielsen, P. J. (2007): Metabolic effects of interventions to increase exercise in adults with type 2 diabetes. In: *Diabetologia* 50 (5), S. 913–921. DOI: 10.1007/s00125-007-0625-0.
- Connelly, J.; Kirk, A.; Masthoff, J.; MacRury, S. (2013): The use of technology to promote physical activity in Type 2 diabetes management: a systematic review. In: *Diabetic medicine : a journal of the British Diabetic Association* 30 (12), S. 1420–1432. DOI: 10.1111/dme.12289.
- Conraads, V. M.; Beckers, P.; Bosmans, J.; Clerck, L. S. de; Stevens, W. J.; Vrints, C. J.; Brutsaert, D. L. (2002): Combined endurance/resistance training reduces plasma TNF-alpha receptor levels in patients with chronic heart failure and coronary artery disease. In: *European Heart Journal* 23 (23), S. 1854–1860. DOI: 10.1053/euhj.2002.3239.
- Cook, Nancy R. (1995): Implications of Small Reductions in Diastolic Blood Pressure for Primary Prevention. In: *Arch Intern Med* 155 (7), S. 701. DOI: 10.1001/archinte.1995.00430070053006.
- Cooney (2013): Exercise for depression. In: *Cochrane Database Syst Rev* 9, CD004366.
- Cornelissen, Véronique A.; Buys, Roselien; Smart, Neil A. (2013): Endurance exercise beneficially affects ambulatory blood pressure: a systematic review and meta-analysis. In: *Journal of Hypertension* 31 (4), S. 639–648. DOI: 10.1097/HJH.0b013e32835ca964.
- Cornelissen, Véronique A.; Fagard, Robert H. (2005): Effect of resistance training on resting blood pressure: a meta-analysis of randomized controlled trials. In: *Journal of Hypertension* 23 (2), S. 251–259. DOI: 10.1097/00004872-200502000-00003.
- Cornelissen, Véronique A.; Fagard, Robert H.; Coeckelberghs, Ellen; Vanhees, Luc (2011): Impact of resistance training on blood pressure and other cardiovascular risk factors: a meta-analysis of randomized, controlled trials. In: *Hypertension* 58 (5), S. 950–958. DOI: 10.1161/HYPERTENSIONAHA.111.177071.
- Cornelissen, Veronique A.; Smart, Neil A. (2013): Exercise training for blood pressure: a systematic review and meta-analysis. In: *Journal of the American Heart Association* 2 (1), e004473. DOI: 10.1161/JAHA.112.004473.
- Courneya, K. S.; Friedenreich, C. M. (1999): Physical exercise and quality of life following cancer diagnosis: a literature review. In: *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine* 21 (2), S. 171–179. DOI: 10.1007/BF02908298.
- Courneya, K. S.; Mackey, J. R.; Jones, L. W. (2000): Coping with cancer: can exercise help? In: *The Physician and Sportsmedicine* 28 (5), S. 49–73. DOI: 10.3810/psm.2000.05.896.
- Crawford, D. A.; Jeffery, R. W.; French, S. A. (1999): Television viewing, physical inactivity and obesity. In: *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 23 (4), S. 437–440. DOI: 10.1038/sj.ijo.0800845.
- Creasy, T. S.; McMillan, P. J.; Fletcher, E.W.L.; Collin, J.; Morris, P. J. (1990): Is percutaneous transluminal angioplasty better than exercise for claudication?—Preliminary results from a prospective randomised trial. In: *European Journal of Vascular Surgery* 4 (2), S. 135–140. DOI: 10.1016/S0950-821X(05)80427-X.
- Creatas, George; Deligeoroglou, Efthimios (2007): Polycystic ovarian syndrome in adolescents. In: *Current Opinion in Obstetrics and Gynecology* 19 (5), S. 420–426. DOI: 10.1097/GCO.0b013e3282efdc54.
- Crouse, S. F.; O'Brien, B. C.; Grandjean, P. W.; Lowe, R. C.; Rohack, J. J.; Green, J. S.; Tolson, H. (1997): Training intensity, blood lipids, and apolipoproteins in men with high cholesterol. In: *Journal of applied physiology (Bethesda, Md. : 1985)* 82 (1), S. 270–277. DOI: 10.1152/jappl.1997.82.1.270.
- Cuenca-García, M.; Jago, R.; Shield, J. P. H.; Burren, C. P. (2012): How does physical activity and fitness influence glycaemic control in young people with Type 1 diabetes? In: *Diabetic medicine : a journal of the British Diabetic Association* 29 (10), e369–76. DOI: 10.1111/j.1464-5491.2012.03740.x.
- Cummings, S. R.; Nevitt, M. C.; Browner, W. S.; Stone, K.; Fox, K. M.; Ensrud, K. E. et al. (1995): Risk factors for hip fracture in white women. Study of Osteoporotic Fractures Research Group. In: *The New England journal of medicine* 332 (12), S. 767–773. DOI: 10.1056/NEJM199503233321202.
- Dagfinrud (2008): Physiotherapy interventions for ankylosing spondylitis. In: *Cochrane Database Syst Rev*, CD002822.
- Dale (1990): Weight maintenance and resting metabolic rate 18–40 months after a diet/exercise treatment. In: *Int J Obes* 14, S. 347.
- Daley, A. J.; Foster, L.; Long, G.; Palmer, C.; Robinson, O.; Walmsley, H.; Ward, R. (2015): The effectiveness of exercise for the prevention and treatment of antenatal depression: systematic review with meta-analysis. In: *BJOG : an international journal of obstetrics and gynaecology* 122 (1), S. 57–62. DOI: 10.1111/1471-0528.12909.
- Dalgas, Ulrik; Stenager, Egon (2012): Exercise and disease progression in multiple sclerosis: can exercise slow down the progression of multiple sclerosis? In: *Therapeutic Advances in Neurological Disorders* 5 (2), S. 81–95. DOI: 10.1177/1756285611430719.
- Dalgas, U.; Stenager, E.; Jakobsen, J.; Petersen, T.; Hansen, H. J.; Knudsen, C. et al. (2009): Resistance training improves muscle strength and functional capacity in multiple sclerosis. In: *Neurology* 73 (18), S. 1478–1484. DOI: 10.1212/WNL.0b013e3181bf98b4.

- Dalgas, U.; Stenager, E.; Jakobsen, J.; Petersen, T.; Hansen, H. J.; Knudsen, C. et al. (2010): Fatigue, mood and quality of life improve in MS patients after progressive resistance training. In: *Multiple Sclerosis Journal* 16 (4), S. 480–490. DOI: 10.1177/1352458509360040.
- Davidson, Lance E.; Hudson, Robert; Kilpatrick, Katherine; Kuk, Jennifer L.; McMillan, Kathleen; Janiszewski, Peter M. et al. (2009): Effects of exercise modality on insulin resistance and functional limitation in older adults: a randomized controlled trial. In: *Arch Intern Med* 169 (2), S. 122–131. DOI: 10.1001/archinternmed.2008.558.
- Davies (2010): Exercise based rehabilitation for heart failure. In: *Cochrane Database Syst Rev*, CD003331.
- Davies, Edward J.; Moxham, Tiffany; Rees, Karen; Singh, Sally; Coats, Andrew J. S.; Ebrahim, Shah et al. (2010): Exercise training for systolic heart failure: Cochrane systematic review and meta-analysis. In: *European Journal of Heart Failure* 12 (7), S. 706–715. DOI: 10.1093/eurjhf/hfq056.
- Davis, P. B.; Drumm, M.; Konstan, M. W. (1996): Cystic fibrosis. In: *Am J Respir Crit Care Med* 154 (5), S. 1229–1256. DOI: 10.1164/ajrccm.154.5.8912731.
- Day, Lesley; Fildes, Brian; Gordon, Ian; Fitzharris, Michael; Flamer, Harold; Lord, Stephen (2002): Randomised factorial trial of falls prevention among older people living in their own homes. In: *BMJ* 325 (7356), S. 128. DOI: 10.1136/bmj.325.7356.128.
- Dela, F.; Handberg, A.; Mikines, K. J.; Vinten, J.; Galbo, H. (1993): GLUT 4 and insulin receptor binding and kinase activity in trained human muscle. In: *The Journal of Physiology* 469 (1), S. 615–624. DOI: 10.1113/jphysiol.1993.sp019833.
- Dela, F.; Larsen, J. J.; Mikines, K. J.; Ploug, T.; Petersen, L. N.; Galbo, H. (1995): Insulin-stimulated muscle glucose clearance in patients with NIDDM. Effects of one-legged physical training. In: *Diabetes* 44 (9), S. 1010–1020. DOI: 10.2337/diab.44.9.1010.
- Dela, F.; Ploug, T.; Handberg, A.; Petersen, L. N.; Larsen, J. J.; Mikines, K. J.; Galbo, H. (1994): Physical training increases muscle GLUT4 protein and mRNA in patients with NIDDM. In: *Diabetes* 43 (7), S. 862–865. DOI: 10.2337/diab.43.7.862.
- Demopoulos, Laura; Bijou, Rachel; Fergus, Icilma; Jones, Margaret; Strom, Joel; LeJemtel, Thierry H. (1997): Exercise Training in Patients With Severe Congestive Heart Failure: Enhancing Peak Aerobic Capacity While Minimizing the Increase in Ventricular Wall Stress. In: *Journal of the American College of Cardiology* 29 (3), S. 597–603. DOI: 10.1016/S0735-1097(96)00526-8.
- DePue, J. D.; Clark, M. M.; Ruggiero, L.; Medeiros, M. L.; Pera, V. (1995): Maintenance of weight loss: a needs assessment. In: *Obesity Research* 3 (3), S. 241–248. DOI: 10.1002/j.1550-8528.1995.tb00144.x.
- Derby, Carol A.; Mohr, Beth A.; Goldstein, Irwin; Feldman, Henry A.; Johannes, Catherine B.; McKinlay, John B. (2000): Modifiable risk factors and erectile dysfunction: can lifestyle changes modify risk? In: *Urology* 56 (2), S. 302–306. DOI: 10.1016/S0090-4295(00)00614-2.
- Dereli, Elif E.; Yaliman, Ayse (2010): Comparison of the effects of a physiotherapist-supervised exercise programme and a self-supervised exercise programme on quality of life in patients with Parkinson's disease. In: *Clinical Rehabilitation* 24 (4), S. 352–362. DOI: 10.1177/0269215509358933.
- Devore, Elizabeth E.; Kang, Jae Hee; Okereke, Olivia; Grodstein, Francine (2009): Physical activity levels and cognition in women with type 2 diabetes. In: *American Journal of Epidemiology* 170 (8), S. 1040–1047. DOI: 10.1093/aje/kwp224.
- Di (1993): Exercise therapy of non-insulin dependent diabetes mellitus a report of 10 year studies. The efficacy of exercise therapy. In: *Chin Med J (Engl)* 106, S. 757.
- Dietz (1981): Rehabilitation oncology.
- Dimeo, Fernando C.; Stieglitz, Rolf-Dieter; Novelli-Fischer, Ulrike; Fetscher, Sebastian; Keul, Joseph (1999): Effects of physical activity on the fatigue and psychologic status of cancer patients during chemotherapy. In: *Cancer* 85 (10), S. 2273–2277. DOI: 10.1002/(SICI)1097-0142(19990515)85:10<2273::AID-CNCR24>3.0.CO;2-B.
- Donnellan, C.; Hickey, A.; Hevey, D.; O'Neill, D. (2010): Effect of mood symptoms on recovery one year after stroke. In: *International Journal of Geriatric Psychiatry* 25 (12), S. 1288–1295. DOI: 10.1002/gps.2482.
- DONNELLY, J.; SMITH, B.; JACOBSEN, D.; KIRK, E.; DUBOSE, K.; HYDER, M. et al. (2004): The role of exercise for weight loss and maintenance. In: *Best Practice & Research Clinical Gastroenterology* 18 (6), S. 1009–1029. DOI: 10.1016/S1521-6918(04)00083-6.
- Dubach, Paul; Myers, Jonathan; Dziekan, Gerald; Goebbels, Ute; Reinhart, Walter; Muller, Peter et al. (1997): Effect of High Intensity Exercise Training on Central Hemodynamic Responses to Exercise in Men With Reduced Left Ventricular Function. In: *Journal of the American College of Cardiology* 29 (7), S. 1591–1598. DOI: 10.1016/S0735-1097(97)82540-5.
- Duijts, Saskia F. A.; Faber, Mizja M.; Oldenburg, Hester S. A.; van Beurden, Marc; Aaronson, Neil K. (2011): Effectiveness of behavioral techniques and physical exercise on psychosocial functioning and health-related quality of life in breast cancer patients and survivors--a meta-analysis. In: *Psycho-Oncology* 20 (2), S. 115–126. DOI: 10.1002/pon.1728.
- Duncan, B. B.; Schmidt, M. I. (2001): Chronic activation of the innate immune system may underlie the metabolic syndrome. In: *Sao Paulo Medical Journal* 119 (3), S. 122–127. DOI: 10.1590/S1516-31802001000300008.
- Dunstan, D. W.; Mori, T. A.; Puddey, I. B.; Beilin, L. J.; Burke, V.; Morton, A. R.; Stanton, K. G. (1997): The independent and combined effects of aerobic exercise and dietary fish intake on serum lipids and glycemic control in NIDDM. A randomized controlled study. In: *Diabetes Care* 20 (6), S. 913–921. DOI: 10.2337/diacare.20.6.913.

- Dwyer (2011): The role of exercise in maintaining health in cystic fibrosis. In: *Curr Opin Pulm Med* 17, S. 455.
- Dwyer, Tiffany J.; Alison, Jennifer A.; McKeough, Zoe J.; Daviskas, Evangelia; Bye, Peter T. P. (2011): Effects of exercise on respiratory flow and sputum properties in patients with cystic fibrosis. In: *Chest* 139 (4), S. 870–877. DOI: 10.1378/chest.10-1158.
- Earhart (2013): Parkinson disease and exercise. In: *Compr Physiol* 3, S. 833.
- Ebeling, P.; Bourey, R.; Koranyi, L.; Tuominen, J. A.; Groop, L. C.; Henriksson, J. et al. (1993): Mechanism of enhanced insulin sensitivity in athletes. Increased blood flow, muscle glucose transport protein (GLUT-4) concentration, and glycogen synthase activity. In: *Journal of Clinical Investigation* 92 (4), S. 1623–1631. DOI: 10.1172/JCI116747.
- Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34) (1998). In: *The Lancet* 352 (9131), S. 854–865.
- Eid, A. A.; Ionescu, A. A.; Nixon, L. S.; Lewis-Jenkins, V.; Matthews, S. B.; Griffiths, T. L.; Shale, D. J. (2001): Inflammatory response and body composition in chronic obstructive pulmonary disease. In: *Am J Respir Crit Care Med* 164 (8 Pt 1), S. 1414–1418. DOI: 10.1164/ajrccm.164.8.2008109.
- Eklom, B.; Lövgren, O.; Alderin, M.; Fridström, M.; Sätterström, G. (1974): Physical performance in patients with rheumatoid arthritis. In: *Scandinavian Journal of Rheumatology* 3 (3), S. 121–125. DOI: 10.3109/03009747409097136.
- Ekdahl, C.; Broman, G. (1992): Muscle strength, endurance, and aerobic capacity in rheumatoid arthritis: a comparative study with healthy subjects. In: *Annals of the Rheumatic Diseases* 51 (1), S. 35–40. DOI: 10.1136/ard.51.1.35.
- Ekeland (2004): Exercise to improve self-esteem in children and young people. In: *Cochrane Database Syst Rev*, CD003683.
- Eleftheriadou, M.; Michala, L.; Stefanidis, K.; Iliadis, I.; Lykeridou, A.; Antsaklis, A. (2012): Exercise and sedentary habits among adolescents with PCOS. In: *Journal of Pediatric and Adolescent Gynecology* 25 (3), S. 172–174. DOI: 10.1016/j.jpag.2011.11.009.
- Eliassen, A. Heather; Hankinson, Susan E.; Rosner, Bernard; Holmes, Michelle D.; Willett, Walter C. (2010): Physical activity and risk of breast cancer among postmenopausal women. In: *Arch Intern Med* 170 (19), S. 1758–1764. DOI: 10.1001/archinternmed.2010.363.
- Ellekjaer, H.; Holmen, J.; Ellekjaer, E.; Vatten, L. (2000): Physical activity and stroke mortality in women. Ten-year follow-up of the Nord-Trøndelag health survey, 1984–1986. In: *Stroke* 31 (1), S. 14–18. DOI: 10.1161/01.STR.31.1.14.
- Elliott, T. G.; Cockcroft, J. R.; Groop, P. H.; Viberti, G. C.; Ritter, J. M. (1993): Inhibition of nitric oxide synthesis in forearm vasculature of insulin-dependent diabetic patients: blunted vasoconstriction in patients with microalbuminuria. In: *Clinical science (London, England : 1979)* 85 (6), S. 687–693. DOI: 10.1042/cs0850687.
- Ellis, Terry; Latham, Nancy K.; DeAngelis, Tamara R.; Thomas, Cathi A.; Saint-Hilaire, Marie; Bickmore, Timothy W. (2013): Feasibility of a virtual exercise coach to promote walking in community-dwelling persons with Parkinson disease. In: *American Journal of Physical Medicine & Rehabilitation* 92 (6), 472-81; quiz 482-5. DOI: 10.1097/PHM.0b013e31828cd466.
- Elwood, Peter; Galante, Julieta; Pickering, Janet; Palmer, Stephen; Bayer, Antony; Ben-Shlomo, Yoav et al. (2013): Healthy lifestyles reduce the incidence of chronic diseases and dementia: evidence from the Caerphilly cohort study. In: *PLoS ONE* 8 (12), e81877. DOI: 10.1371/journal.pone.0081877.
- Emtner, Margareta; Finne, Merja; Stålenheim, Gunnemar (1998): A 3-year follow-up of asthmatic patients participating in a 10-week rehabilitation program with emphasis on physical training. In: *Archives of Physical Medicine and Rehabilitation* 79 (5), S. 539–544. DOI: 10.1016/S0003-9993(98)90070-3.
- Emtner, M.; Herala, M.; Stålenheim, G. (1996): High-intensity physical training in adults with asthma. A 10-week rehabilitation program. In: *Chest* 109 (2), S. 323–330. DOI: 10.1378/chest.109.2.323.
- Ensari, Ipek; Motl, Robert W.; Pilutti, Lara A. (2014): Exercise training improves depressive symptoms in people with multiple sclerosis: results of a meta-analysis. In: *Journal of Psychosomatic Research* 76 (6), S. 465–471. DOI: 10.1016/j.jpsychores.2014.03.014.
- Erickson, Kirk I.; Voss, Michelle W.; Prakash, Ruchika Shaurya; Basak, Chandramallika; Szabo, Amanda; Chaddock, Laura et al. (2011): Exercise training increases size of hippocampus and improves memory. In: *Proceedings of the National Academy of Sciences* 108 (7), S. 3017–3022. DOI: 10.1073/pnas.1015950108.
- Eriksson, K. F.; Lindgärde, F. (1998): No excess 12-year mortality in men with impaired glucose tolerance who participated in the Malmö Preventive Trial with diet and exercise. In: *Diabetologia* 41 (9), S. 1010–1016. DOI: 10.1007/s001250051024.
- Eriksson, J.; Tuominen, J.; Valle, T.; Sundberg, S.; Sovijärvi, A.; Lindholm, H. et al. (1998): Aerobic endurance exercise or circuit-type resistance training for individuals with impaired glucose tolerance? In: *Hormone and metabolic research = Hormon- und Stoffwechselforschung = Hormones et métabolisme* 30 (1), S. 37–41. DOI: 10.1055/s-2007-978828.
- Esler, M. (2001): The sympathetic neurobiology of essential hypertension: disparate influences of obesity, stress, and noradrenaline transporter dysfunction? In: *American Journal of Hypertension* 14 (11), S139–S146. DOI: 10.1016/S0895-7061(01)02081-7.



- Esposito, Katherine; Giugliano, Francesco; Di Palo, Carmen; Giugliano, Giovanni; Marfella, Raffaele; D'Andrea, Francesco et al. (2004): Effect of lifestyle changes on erectile dysfunction in obese men: a randomized controlled trial. In: *JAMA* 291 (24), S. 2978–2984. DOI: 10.1001/jama.291.24.2978.
- Ewbank, P. P.; Darga, L. L.; Lucas, C. P. (1995): Physical activity as a predictor of weight maintenance in previously obese subjects. In: *Obesity Research* 3 (3), S. 257–263. DOI: 10.1002/j.1550-8528.1995.tb00146.x.
- Fagard, Robert H. (2005): Physical activity, physical fitness and the incidence of hypertension. In: *Journal of Hypertension* 23 (2), S. 265–267. DOI: 10.1097/00004872-200502000-00005.
- Fagard, Robert H.; Cornelissen, Véronique A. (2007): Effect of exercise on blood pressure control in hypertensive patients. In: *European Journal of Cardiovascular Prevention & Rehabilitation* 14 (1), S. 12–17. DOI: 10.1097/HJR.0b013e3280128bbb.
- Fairfield, Kathleen M.; Fletcher, Robert H. (2002): Vitamins for chronic disease prevention in adults: scientific review. In: *JAMA* 287 (23), S. 3116–3126. DOI: 10.1001/jama.287.23.3116.
- Falloon, I. R.; Talbot, R. E. (1981): Persistent auditory hallucinations: coping mechanisms and implications for management. In: *Psychological Medicine* 11 (2), S. 329–339. DOI: 10.1017/S0033291700052144.
- Farmer, M. E.; Harris, T.; Madans, J. H.; Wallace, R. B.; Cornoni-Huntley, J.; White, L. R. (1989): Anthropometric indicators and hip fracture. The NHANES I epidemiologic follow-up study. In: *Journal of the American Geriatrics Society* 37 (1), S. 9–16. DOI: 10.1111/j.1532-5415.1989.tb01562.x.
- FARRELL, STEPHEN W.; FINLEY, CARRIE E.; GRUNDY, SCOTT M. (2012): Cardiorespiratory fitness, LDL cholesterol, and CHD mortality in men. In: *Medicine and science in sports and exercise* 44 (11), S. 2132–2137. DOI: 10.1249/MSS.0b013e31826524be.
- Faulkner, Guy; Cohn, Tony; Remington, Gary (2006): Validation of a physical activity assessment tool for individuals with schizophrenia. In: *Schizophrenia Research* 82 (2–3), S. 225–231. DOI: 10.1016/j.schres.2005.10.020.
- FEBBRAIO, MARK A.; PEDERSEN, BENTE KLARLUND (2002): Muscle-derived interleukin-6: mechanisms for activation and possible biological roles. In: *The FASEB Journal* 16 (11), S. 1335–1347. DOI: 10.1096/fj.01-0876rev.
- Felson, D. T.; Naimark, A.; Anderson, J.; Kazis, L.; Castelli, W.; Meenan, R. F. (1987): The prevalence of knee osteoarthritis in the elderly. The Framingham Osteoarthritis Study. In: *Arthritis and rheumatism* 30 (8), S. 914–918. DOI: 10.1002/art.1780300811.
- Fillit, Howard; Nash, David T.; Rundek, Tatjana; Zuckerman, Andrea (2008): Cardiovascular risk factors and dementia. In: *The American Journal of Geriatric Pharmacotherapy* 6 (2), S. 100–118. DOI: 10.1016/j.amjopharm.2008.06.004.
- Firth, J.; Cotter, J.; Elliott, R.; French, P.; Yung, A. R. (2015): A systematic review and meta-analysis of exercise interventions in schizophrenia patients. In: *Psychological Medicine* 45 (7), S. 1343–1361. DOI: 10.1017/S0033291714003110.
- Fitzgerald, G. Kelley; Piva, Sara R.; Irrgang, James J.; Bouzubar, Fawzi; Starz, Terence W. (2004): Quadriceps activation failure as a moderator of the relationship between quadriceps strength and physical function in individuals with knee osteoarthritis. In: *Arthritis and rheumatism* 51 (1), S. 40–48. DOI: 10.1002/art.20084.
- Fleshner, Monika (2005): Physical activity and stress resistance: sympathetic nervous system adaptations prevent stress-induced immunosuppression. In: *Exercise and Sport Sciences Reviews* 33 (3), S. 120–126. DOI: 10.1097/00003677-200507000-00004.
- Flynn, Kathryn E.; Piña, Ileana L.; Whellan, David J.; Lin, Li; Blumenthal, James A.; Ellis, Stephen J. et al. (2009): Effects of exercise training on health status in patients with chronic heart failure: HF-ACTION randomized controlled trial. In: *JAMA* 301 (14), S. 1451–1459. DOI: 10.1001/jama.2009.457.
- Flynn, T. J.; Walsh, M. F. (1993): Thirty-month evaluation of a popular very-low-calorie diet program. In: *Archives of Family Medicine* 2 (10), S. 1042–1048. DOI: 10.1001/archfami.2.10.1042.
- Fogelholm, M. (2010): Physical activity, fitness and fatness: relations to mortality, morbidity and disease risk factors. A systematic review. In: *Obesity Reviews* 11 (3), S. 202–221. DOI: 10.1111/j.1467-789X.2009.00653.x.
- Fogelholm, M.; Kukkonen-Harjula, K. (2000): Does physical activity prevent weight gain--a systematic review. In: *Obesity reviews : an official journal of the International Association for the Study of Obesity* 1 (2), S. 95–111. DOI: 10.1046/j.1467-789x.2000.00016.x.
- Fogelholm, M.; Kukkonen-Harjula, K.; Nenonen, A.; Pasanen, M. (2000): Effects of walking training on weight maintenance after a very-low-energy diet in premenopausal obese women: a randomized controlled trial. In: *Archives of Internal Medicine* 160 (14), S. 2177–2184. DOI: 10.1001/archinte.160.14.2177.
- Forbes (2015): Exercise programs for people with dementia. In: *Cochrane Database Syst Rev* 4, CD006489.
- Ford, Earl S. (2005): Prevalence of the metabolic syndrome defined by the International Diabetes Federation among adults in the U.S. In: *Diabetes Care* 28 (11), S. 2745–2749. DOI: 10.2337/diacare.28.11.2745.
- Førde, O. H.; Thelle, D. S.; Arnesen, E.; Mjø, O. D. (1986): Distribution of high density lipoprotein cholesterol according to relative body weight, cigarette smoking and leisure time physical activity. The Cardiovascular Disease Study in Finnmark 1977. In: *Acta Medica Scandinavica* 219 (2), S. 167–171. DOI: 10.1111/j.0954-6820.1986.tb03294.x.
- Forouhi, N. G.; Sattar, N.; McKeigue, P. M. (2001): Relation of C-reactive protein to body fat distribution and features of the metabolic syndrome in Europeans and South Asians. In: *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 25 (9), S. 1327–1331. DOI: 10.1038/sj.ijo.0801723.

- Fox (1998): Fox's physiological basis for exercise physiology.
- Fransen (2015): Exercise for osteoarthritis of the knee. In: *Cochrane Database Syst Rev* 1, CD004376.
- Frazzitta, Giuseppe; Balbi, Pietro; Maestri, Roberto; Bertotti, Gabriella; Boveri, Natalia; Pezzoli, Gianni (2013): The beneficial role of intensive exercise on Parkinson disease progression. In: *American Journal of Physical Medicine & Rehabilitation* 92 (6), S. 523–532. DOI: 10.1097/PHM.0b013e31828cd254.
- Fuchsjaeger-Mayrl, Gabriele; Pleiner, Johannes; Wiesinger, Günther F.; Sieder, Anna E.; Quittan, Michael; Nuhr, Martin J. et al. (2002): Exercise training improves vascular endothelial function in patients with type 1 diabetes. In: *Diabetes Care* 25 (10), S. 1795–1801. DOI: 10.2337/diacare.25.10.1795.
- Gabriel, Sherine E. (2008): Cardiovascular morbidity and mortality in rheumatoid arthritis. In: *The American Journal of Medicine* 121 (10 Suppl 1), S9-14. DOI: 10.1016/j.amjmed.2008.06.011.
- Gaede, Peter; Vedel, Pernille; Larsen, Nicolai; Jensen, Gunnar V. H.; Parving, Hans-Henrik; Pedersen, Oluf (2003): Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. In: *New England Journal of Medicine* 348 (5), S. 383–393. DOI: 10.1056/NEJMoa021778.
- Galipeau, Denise M.; Yao, Linfu; McNeill, John H. (2002): Relationship among hyperinsulinemia, insulin resistance, and hypertension is dependent on sex. In: *American journal of physiology. Heart and circulatory physiology* 283 (2), H562-7. DOI: 10.1152/ajpheart.00238.2002.
- GALPER, DANIEL I.; TRIVEDI, MADHUKAR H.; BARLOW, CAROLYN E.; DUNN, ANDREA L.; KAMPERT, JAMES B. (2006): Inverse association between physical inactivity and mental health in men and women. In: *Medicine & Science in Sports & Exercise* 38 (1), S. 173–178. DOI: 10.1249/01.mss.0000180883.32116.28.
- García-Hermoso, A.; Saavedra, J. M.; Escalante, Y. (2013): Effects of exercise on resting blood pressure in obese children: a meta-analysis of randomized controlled trials. In: *Obesity Reviews* 14 (11), S. 919–928. DOI: 10.1111/obr.12054.
- Garfinkel, S. K.; Kesten, S.; Chapman, K. R.; Rebuck, A. S. (1992): Physiologic and nonphysiologic determinants of aerobic fitness in mild to moderate asthma. In: *American Review of Respiratory Disease* 145 (4 Pt 1), S. 741–745. DOI: 10.1164/ajrccm/145.4\_Pt\_1.741.
- George, Stephanie M.; Irwin, Melinda L.; Smith, Ashley W.; Neuhaus, Marian L.; Reedy, Jill; McTiernan, Anne et al. (2011): Postdiagnosis diet quality, the combination of diet quality and recreational physical activity, and prognosis after early-stage breast cancer. In: *Cancer Causes & Control* 22 (4), S. 589–598. DOI: 10.1007/s10552-011-9732-9.
- Georgiades, A.; Sherwood, A.; Gullette, E. C.; Babyak, M. A.; Hinderliter, A.; Waugh, R. et al. (2000): Effects of exercise and weight loss on mental stress-induced cardiovascular responses in individuals with high blood pressure. In: *Hypertension* 36 (2), S. 171–176. DOI: 10.1161/01.HYP.36.2.171.
- Giannopoulou, I.; Ploutz-Snyder, L. L.; Carhart, R.; Weinstock, R. S.; Fernhall, B.; Goulopoulou, S.; Kanaley, J. A. (2005): Exercise is required for visceral fat loss in postmenopausal women with type 2 diabetes. In: *The Journal of clinical endocrinology and metabolism* 90 (3), S. 1511–1518. DOI: 10.1210/jc.2004-1782.
- Giannuzzi, P.; Mezzani, A.; Saner, H.; Björnstad, H.; Fioretti, P.; Mendes, M. et al. (2003): Physical activity for primary and secondary prevention. Position paper of the Working Group on Cardiac Rehabilitation and Exercise Physiology of the European Society of Cardiology. In: *European Journal of Cardiovascular Prevention & Rehabilitation* 10 (5), S. 319–327. DOI: 10.1097/01.hjr.0000086303.28200.50.
- Gielen, Stephan; Adams, Volker; Möbius-Winkler, Sven; Linke, Axel; Erbs, Sandra; Yu, Jiangtao et al. (2003): Anti-inflammatory effects of exercise training in the skeletal muscle of patients with chronic heart failure. In: *Journal of the American College of Cardiology* 42 (5), S. 861–868. DOI: 10.1016/S0735-1097(03)00848-9.
- Gillespie (2001): Interventions for preventing falls in elderly people. In: *Cochrane Database Syst Rev*, CD000340.
- Glenthøj, Birte Y.; Hemmingsen, Ralf (1997): Dopaminergic sensitization: Implications for the pathogenesis of schizophrenia. In: *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 21 (1), S. 23–46. DOI: 10.1016/S0278-5846(96)00158-3.
- Goede, C. J. de; Keus, S. H.; Kwakkel, G.; Wagenaar, R. C. (2001): The effects of physical therapy in Parkinson's disease: a research synthesis. In: *Archives of Physical Medicine and Rehabilitation* 82 (4), S. 509–515. DOI: 10.1053/apmr.2001.22352.
- Gokce, Noyan; Vita, Joseph A.; Bader, David S.; Sherman, Debra L.; Hunter, Liza M.; Holbrook, Monika et al. (2002): Effect of exercise on upper and lower extremity endothelial function in patients with coronary artery disease. In: *The American Journal of Cardiology* 90 (2), S. 124–127. DOI: 10.1016/S0002-9149(02)02433-5.
- Goldbourt, U.; Yaari, S.; Medalie, J. H. (1993): Factors predictive of long-term coronary heart disease mortality among 10,059 male Israeli civil servants and municipal employees. A 23-year mortality follow-up in the Israeli Ischemic Heart Disease Study. In: *Cardiology* 82 (2-3), S. 100–121. DOI: 10.1159/000175862.
- Gommans, Lindy N. M.; Fokkenrood, Hugo J. P.; van Dalen, Hendrika C. W.; Scheltinga, Marc R. M.; Teijink, Joep A. W.; Peters, Ron J. G. (2015): Safety of supervised exercise therapy in patients with intermittent claudication. In: *Journal of Vascular Surgery* 61 (2), 512-518.e2. DOI: 10.1016/j.jvs.2014.08.070.

- Goodwin, Victoria A.; Richards, Suzanne H.; Taylor, Rod S.; Taylor, Adrian H.; Campbell, John L. (2008): The effectiveness of exercise interventions for people with Parkinson's disease: a systematic review and meta-analysis. In: *Movement Disorders* 23 (5), S. 631–640. DOI: 10.1002/mds.21922.
- Gorczyński (2010): Exercise therapy for schizophrenia. In: *Cochrane Database Syst Rev*, CD004412.
- Gosselink, Rik; Kovacs, Ludwig; Ketelaer, Pierre; Carton, Herwig; Decramer, Marc (2000): Respiratory muscle weakness and respiratory muscle training in severely disabled multiple sclerosis patients. In: *Archives of Physical Medicine and Rehabilitation* 81 (6), S. 747–751. DOI: 10.1016/S0003-9993(00)90105-9.
- Green, R. H.; Singh, S. J.; Williams, J.; Morgan, M. D. (2001): A randomised controlled trial of four weeks versus seven weeks of pulmonary rehabilitation in chronic obstructive pulmonary disease. In: *Thorax* 56 (2), S. 143–145. DOI: 10.1136/thorax.56.2.143.
- Green, John S.; Stanforth, Philip R.; Rankinen, Tuomo; Leon, Arthur S.; Rao Dc, D. c.; Skinner, James S. et al. (2004): The effects of exercise training on abdominal visceral fat, body composition, and indicators of the metabolic syndrome in postmenopausal women with and without estrogen replacement therapy: the HERITAGE family study. In: *Metabolism* 53 (9), S. 1192–1196. DOI: 10.1016/j.metabol.2004.04.008.
- Griffiths, T. L.; Burr, M. L.; Campbell, I. A.; Lewis-Jenkins, V.; Mullins, J.; Shiels, K. et al. (2000): Results at 1 year of outpatient multidisciplinary pulmonary rehabilitation: a randomised controlled trial. In: *The Lancet* 355 (9201), S. 362–368. DOI: 10.1016/S0140-6736(99)07042-7.
- Griffiths, T. L.; Phillips, C. J.; Davies, S.; Burr, M. L.; Campbell, I. A. (2001): Cost effectiveness of an outpatient multidisciplinary pulmonary rehabilitation programme. In: *Thorax* 56 (10), S. 779–784. DOI: 10.1136/thorax.56.10.779.
- Grodstein, Francine (1996): Three-Year Follow-up of Participants in a Commercial Weight Loss Program. In: *Archives of Internal Medicine* 156 (12), S. 1302. DOI: 10.1001/archinte.1996.00440110068009.
- Gruber, Wolfgang; Orenstein, David M.; Braumann, Klaus M.; Beneke, Ralph (2014): Interval exercise training in cystic fibrosis -- effects on exercise capacity in severely affected adults. In: *Journal of cystic fibrosis : official journal of the European Cystic Fibrosis Society* 13 (1), S. 86–91. DOI: 10.1016/j.jcf.2013.06.005.
- Guelfi, Kym J.; Jones, Timothy W.; Fournier, Paul A. (2007): New insights into managing the risk of hypoglycaemia associated with intermittent high-intensity exercise in individuals with type 1 diabetes mellitus: implications for existing guidelines. In: *Sports Medicine* 37 (11), S. 937–946. DOI: 10.2165/00007256-200737110-00002.
- Guelfi, K. J.; Ratnam, N.; Smythe, G. A.; Jones, T. W.; Fournier, P. A. (2007): Effect of intermittent high-intensity compared with continuous moderate exercise on glucose production and utilization in individuals with type 1 diabetes. In: *American journal of physiology. Endocrinology and metabolism* 292 (3), E865-70. DOI: 10.1152/ajpendo.00533.2006.
- Gueyffier, F.; Boutitie, F.; Boissel, J. P.; Pocock, S.; Coope, J.; Cutler, J. et al. (1997): Effect of antihypertensive drug treatment on cardiovascular outcomes in women and men. A meta-analysis of individual patient data from randomized, controlled trials. The INDANA Investigators. In: *Annals of Internal Medicine* 126 (10), S. 761–767. DOI: 10.7326/0003-4819-126-10-199705150-00002.
- Gunnarsson (1987): Serum lipid and lipoprotein levels in female type I diabetics: relationships to aerobic capacity and glycaemic control. In: *Diabete Metab* 13, S. 417.
- Guo, S. S.; Zeller, C.; Chumlea, W. C.; Siervogel, R. M. (1999): Aging, body composition, and lifestyle: the Fels Longitudinal Study. In: *The American Journal of Clinical Nutrition* 70 (3), S. 405–411. DOI: 10.1093/ajcn/70.3.405.
- Gustafsson, T.; Bodin, K.; Sylvén, C.; Gordon, A.; Tyini-Lenné, R.; Jansson, E. (2001): Increased expression of VEGF following exercise training in patients with heart failure. In: *European Journal of Clinical Investigation* 31 (4), S. 362–366. DOI: 10.1046/j.1365-2362.2001.00816.x.
- Guzick, David S. (2008): Do cardiovascular risk factors in polycystic ovarian syndrome result in more cardiovascular events? In: *The Journal of clinical endocrinology and metabolism* 93 (4), S. 1170–1171. DOI: 10.1210/jc.2008-0288.
- Haapanen, N.; Miilunpalo, S.; Vuori, I.; Oja, P.; Pasanen, M. (1997): Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women. In: *International Journal of Epidemiology* 26 (4), S. 739–747. DOI: 10.1093/ije/26.4.739.
- Hagen, K. B.; Hilde, G.; Jamtvedt, G.; Winnem, M. F. (2000): The Cochrane review of bed rest for acute low back pain and sciatica. In: *Spine* 25 (22), S. 2932–2939. DOI: 10.1097/00007632-200011150-00016.
- Hakkinen (1999): Dynamic strength training in patients with early rheumatoid arthritis increases muscle strength but not bone mineral density. In: *J Rheumatol* 26, S. 1257.
- Hakkinen (2005): Effects of prolonged combined strength and endurance training on physical fitness, body composition and serum hormones in women with rheumatoid arthritis and in healthy controls. In: *Clin Exp Rheumatol* 23, S. 505.
- HÄKKINEN, A.; HANNONEN, P.; HÄKKINEN, K. (1995): Muscle strength in healthy people and in patients suffering from recent-onset inflammatory arthritis. In: *British journal of rheumatology* 34 (4), S. 355–360. DOI: 10.1093/rheumatology/34.4.355.
- HÄKKINEN, A.; HANNONEN, P.; Nyman, K.; Lyyski, T.; HÄKKINEN, K. (2003): Effects of concurrent strength and endurance training in women with early or longstanding rheumatoid arthritis: comparison with healthy subjects. In: *Arthritis and rheumatism* 49 (6), S. 789–797. DOI: 10.1002/art.11466.

- Halle, Martin; Berg, Aloys; Garwers, Ulrich; Baumstark, Manfred W.; Knisel, Werner; Grathwohl, Dominik et al. (1999): Influence of 4 weeks' intervention by exercise and diet on low-density lipoprotein subfractions in obese men with type 2 diabetes. In: *Metabolism* 48 (5), S. 641–644. DOI: 10.1016/S0026-0495(99)90064-1.
- Hallgreen, C. E.; Hall, K. D. (2008): Allometric relationship between changes of visceral fat and total fat mass. In: *International Journal of Obesity* 32 (5), S. 845–852. DOI: 10.1038/sj.ijo.0803783.
- Hambrecht, R.; Gielen, S.; Linke, A.; Fiehn, E.; Yu, J.; Walther, C. et al. (2000): Effects of exercise training on left ventricular function and peripheral resistance in patients with chronic heart failure: A randomized trial. In: *JAMA* 283 (23), S. 3095–3101. DOI: 10.1001/jama.283.23.3095.
- Hambrecht, Rainer; Niebauer, Josef; Fiehn, Eduard; Kälberer, Barbara; Offner, Bertram; Hauer, Klaus et al. (1995): Physical training in patients with stable chronic heart failure: Effects on cardiorespiratory fitness and ultrastructural abnormalities of leg muscles. In: *Journal of the American College of Cardiology* 25 (6), S. 1239–1249. DOI: 10.1016/0735-1097(94)00568-B.
- Hamilton, A. L.; Killian, K. J.; Summers, E.; Jones, N. L. (1995): Muscle strength, symptom intensity, and exercise capacity in patients with cardiorespiratory disorders. In: *Am J Respir Crit Care Med* 152 (6 Pt 1), S. 2021–2031. DOI: 10.1164/ajrccm.152.6.8520771.
- Hansen, T. M.; Hansen, G.; Langgaard, A. M.; Rasmussen, J. O. (1993): Longterm physical training in rheumatoid arthritis. A randomized trial with different training programs and blinded observers. In: *Scandinavian Journal of Rheumatology* 22 (3), S. 107–112. DOI: 10.3109/03009749309099253.
- Harkcom, T. M.; Lampman, R. M.; Banwell, B. F.; Castor, C. W. (1985): Therapeutic value of graded aerobic exercise training in rheumatoid arthritis. In: *Arthritis and rheumatism* 28 (1), S. 32–39. DOI: 10.1002/art.1780280106.
- Harrington, Derek; Anker, Stefan D.; Chua, Tuan Peng; Webb-Peploe, Katharine M.; Ponikowski, Piotr P.; Poole-Wilson, Philip A.; Coats, Andrew J.S (1997): Skeletal Muscle Function and Its Relation to Exercise Tolerance in Chronic Heart Failure. In: *Journal of the American College of Cardiology* 30 (7), S. 1758–1764. DOI: 10.1016/S0735-1097(97)00381-1.
- Harrison, Paul J. (2004): The hippocampus in schizophrenia: a review of the neuropathological evidence and its pathophysiological implications. In: *Psychopharmacology* 174 (1), S. 151–162. DOI: 10.1007/s00213-003-1761-y.
- Harrison, Cheryce L.; Lombard, Catherine B.; Moran, Lisa J.; Teede, Helena J. (2011): Exercise therapy in polycystic ovary syndrome: a systematic review. In: *Human Reproduction Update* 17 (2), S. 171–183. DOI: 10.1093/humupd/dmq045.
- Harriss, D. J.; Atkinson, G.; Batterham, A.; George, K.; Cable, N. Tim; Reilly, T. et al. (2009): Lifestyle factors and colorectal cancer risk (2): a systematic review and meta-analysis of associations with leisure-time physical activity. In: *Colorectal Disease* 11 (7), S. 689–701. DOI: 10.1111/j.1463-1318.2009.01767.x.
- Hart, Roger (2007): Polycystic ovarian syndrome--prognosis and treatment outcomes. In: *Current Opinion in Obstetrics and Gynecology* 19 (6), S. 529–535. DOI: 10.1097/GCO.0b013e3282f10e22.
- Hartman, William M.; Stroud, Michael; Sweet, David M.; Saxton, Joan (1993): Long-term maintenance of weight loss following supplemented fasting. In: *Int. J. Eat. Disord.* 14 (1), S. 87–93. DOI: 10.1002/1098-108X(199307)14:1<87::AID-EAT2260140112>3.0.CO;2-2.
- Haus, Gail; Hoerr, Sharon L.; Mavis, Brian; Robison, Jon (1994): Key modifiable factors in weight maintenance: Fat intake, exercise, and weight cycling. In: *Journal of the American Dietetic Association* 94 (4), S. 409–413. DOI: 10.1016/0002-8223(94)90096-5.
- Hawkins, P.; Johnson, L. C.; Nikolettou, D.; Hamnegård, C-H; Sherwood, R.; Polkey, M. I.; Moxham, J. (2002): Proportional assist ventilation as an aid to exercise training in severe chronic obstructive pulmonary disease. In: *Thorax* 57 (10), S. 853–859. DOI: 10.1136/thorax.57.10.853.
- Hayashi, Tomoshige; Boyko, Edward J.; Leonetti, Donna L.; McNeely, Marguerite J.; Newell-Morris, Laura; Kahn, Steven E.; Fujimoto, Wilfred Y. (2004): Visceral adiposity is an independent predictor of incident hypertension in Japanese Americans. In: *Annals of Internal Medicine* 140 (12), S. 992–1000. DOI: 10.7326/0003-4819-140-12-200406150-00008.
- Hayashino, Yasuaki; Jackson, Jeffrey L.; Fukumori, Norio; Nakamura, Fumiaki; Fukuhara, Shunichi (2012): Effects of supervised exercise on lipid profiles and blood pressure control in people with type 2 diabetes mellitus: a meta-analysis of randomized controlled trials. In: *Diabetes Research and Clinical Practice* 98 (3), S. 349–360. DOI: 10.1016/j.diabres.2012.10.004.
- Haykowsky, Mark J.; Liang, Yuanyuan; Pechter, David; Jones, Lee W.; McAlister, Finlay A.; Clark, Alexander M. (2007): A meta-analysis of the effect of exercise training on left ventricular remodeling in heart failure patients: the benefit depends on the type of training performed. In: *Journal of the American College of Cardiology* 49 (24), S. 2329–2336. DOI: 10.1016/j.jacc.2007.02.055.
- Haykowsky, Mark J.; Timmons, Meagan P.; Kruger, Calvin; McNeely, Margaret; Taylor, Dylan A.; Clark, Alexander M. (2013): Meta-analysis of aerobic interval training on exercise capacity and systolic function in patients with heart failure and reduced ejection fractions. In: *The American Journal of Cardiology* 111 (10), S. 1466–1469. DOI: 10.1016/j.amjcard.2013.01.303.
- Heitmann, Berit L.; Frederiksen, Peder (2009): Thigh circumference and risk of heart disease and premature death: prospective cohort study. In: *BMJ* 339 (sep03 2), b3292. DOI: 10.1136/bmj.b3292.
- Helge (2001): High prevalence of eating disorders among elite athletes. Increased risk of amenorrhea and premenopausal osteoporosis. In: *Ugeskr Laeger* 163, S. 3473.

- Helmark, Ida C.; Mikkelsen, Ulla R.; Børglum, Jens; Rothe, Anders; Petersen, Marie C. H.; Andersen, Ove et al. (2010): Exercise increases interleukin-10 levels both intraarticularly and peri-synovially in patients with knee osteoarthritis: a randomized controlled trial. In: *Arthritis Research & Therapy* 12 (4), R126. DOI: 10.1186/ar3064.
- Heran (2011): Exercise-based cardiac rehabilitation for coronary heart disease. In: *Cochrane Database Syst Rev*, CD001800.
- Hernández-Molina, Gabriela; Reichenbach, Stephan; Zhang, Bin; Lavalley, Michael; Felson, David T. (2008): Effect of therapeutic exercise for hip osteoarthritis pain: results of a meta-analysis. In: *Arthritis and rheumatism* 59 (9), S. 1221–1228. DOI: 10.1002/art.24010.
- Herring, Matthew P.; O'Connor, Patrick J.; Dishman, Rodney K. (2010): The effect of exercise training on anxiety symptoms among patients: a systematic review. In: *Arch Intern Med* 170 (4), S. 321–331. DOI: 10.1001/archinternmed.2009.530.
- Hiatt, W. R.; Regensteiner, J. G.; Hargarten, M. E.; Wolfel, E. E.; Brass, E. P. (1990): Benefit of exercise conditioning for patients with peripheral arterial disease. In: *Circulation* 81 (2), S. 602–609. DOI: 10.1161/01.CIR.81.2.602.
- Higashi, Y.; Sasaki, S.; Sasaki, N.; Nakagawa, K.; Ueda, T.; Yoshimizu, A. et al. (1999): Daily aerobic exercise improves reactive hyperemia in patients with essential hypertension. In: *Hypertension* 33 (1 Pt 2), S. 591–597. DOI: 10.1161/01.HYP.33.1.591.
- Higashi, Y.; Sasaki, S.; Kurisu, S.; Yoshimizu, A.; Sasaki, N.; Matsuura, H. et al. (1999): Regular aerobic exercise augments endothelium-dependent vascular relaxation in normotensive as well as hypertensive subjects: role of endothelium-derived nitric oxide. In: *Circulation* 100 (11), S. 1194–1202. DOI: 10.1161/01.CIR.100.11.1194.
- Higgins, Timothy P.; Baker, Matthew D.; Evans, Shelley-Ann; Adams, Rachel A.; Cobbold, Christian (2015): Heterogeneous responses of personalised high intensity interval training on type 2 diabetes mellitus and cardiovascular disease risk in young healthy adults. In: *Clinical Hemorheology and Microcirculation* 59 (4), S. 365–377. DOI: 10.3233/CH-141857.
- Hilde (2002): Advice to stay active as a single treatment for low back pain and sciatica. In: *Cochrane Database Syst Rev*, CD003632.
- Hisham (2013): Epidemiology, pathophysiology, and treatment of hypertension in ischaemic stroke patients. In: *J Stroke Cerebrovasc Dis* 22, e4.
- Hkkinen, Arja; Sokka, Tuulikki; Kotaniemi, Antero; Hannonen, Pekka (2001): A randomized two-year study of the effects of dynamic strength training on muscle strength, disease activity, functional capacity, and bone mineral density in early rheumatoid arthritis. In: *Arthritis & Rheumatism* 44 (3), S. 515–522. DOI: 10.1002/1529-0131(200103)44:3<515::AID-ANR98>3.0.CO;2-5.
- Ho, S. C.; Woo, J.; Sham, A.; Chan, S. G.; Yu, A. L. (2001): A 3-year follow-up study of social, lifestyle and health predictors of cognitive impairment in a Chinese older cohort. In: *International Journal of Epidemiology* 30 (6), S. 1389–1396. DOI: 10.1093/ije/30.6.1389.
- Hoeger, Kathleen M. (2008): Exercise therapy in polycystic ovary syndrome. In: *Seminars in Reproductive Medicine* 26 (1), S. 93–100. DOI: 10.1055/s-2007-992929.
- Hoiberg (1984): Correlates of weight loss in treatment and at follow-up. In: *Int J Obes* 8, S. 457.
- Hoidrup (1997): Risk factors for hip fracture.
- Holden (1992): Long-term follow-up of patients attending a combination very-low calorie diet and behaviour therapy weight loss programme. In: *Int J Obes Relat Metab Disord* 16, S. 605.
- Holmes, Michelle D.; Chen, Wendy Y.; Feskanich, Diane; Kroenke, Candyce H.; Colditz, Graham A. (2005): Physical activity and survival after breast cancer diagnosis. In: *JAMA* 293 (20), S. 2479–2486. DOI: 10.1001/jama.293.20.2479.
- Holmes, H.; Ziemba, J.; Evans, T.; Williams, C. A. (1994): Nursing model of psychoeducation for the seriously mentally ill patient. In: *Issues in Mental Health Nursing* 15 (1), S. 85–104. DOI: 10.3109/01612849409074937.
- Holten, Mads K.; Zacho, Morten; Gaster, Michael; Juel, Carsten; Wojtaszewski, Jørgen F. P.; Dela, Flemming (2004): Strength training increases insulin-mediated glucose uptake, GLUT4 content, and insulin signaling in skeletal muscle in patients with type 2 diabetes. In: *Diabetes* 53 (2), S. 294–305. DOI: 10.2337/diabetes.53.2.294.
- Hongo, M.; Itoi, E.; Sinaki, M.; Miyakoshi, N.; Shimada, Y.; Maekawa, S. et al. (2007): Effect of low-intensity back exercise on quality of life and back extensor strength in patients with osteoporosis: a randomized controlled trial. In: *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 18 (10), S. 1389–1395. DOI: 10.1007/s00198-007-0398-9.
- Hørder (1999): Ondt i ryggen. Forekomst, behandling og forebyggelse i et MTV-perspektiv.
- Hornsby, W. G.; Boggess, K. A.; Lyons, T. J.; Barnwell, W. H.; Lazarchick, J.; Colwell, J. A. (1990): Hemostatic alterations with exercise conditioning in NIDDM. In: *Diabetes Care* 13 (2), S. 87–92. DOI: 10.2337/diacare.13.2.87.
- Hoskins (1975): Physiologic responses to known exercise loads in hemiparetic patients. In: *Arch Phys Med Rehabil* 56, S. 544.
- Houmard, Joseph A.; Tanner, Charles J.; Slentz, Cris A.; Duscha, Brian D.; McCartney, Jennifer S.; Kraus, William E. (2004): Effect of the volume and intensity of exercise training on insulin sensitivity. In: *Journal of applied physiology (Bethesda, Md. : 1985)* 96 (1), S. 101–106. DOI: 10.1152/jappphysiol.00707.2003.

- Houston (2013): Inspiratory muscle training for cystic fibrosis. In: *Cochrane Database Syst Rev* 11, CD006112.
- Howe (2011): Exercise for preventing and treating osteoporosis in postmenopausal women. In: *Cochrane Database Syst Rev* 7, CD000333.
- Hsieh (1987): Isokinetic and isometric testing of knee musculature in patients with rheumatoid arthritis with mild knee involvement. In: *Arch Phys Med Rehabil* 68, S. 294.
- Hu, Gang; Jousilahti, Pekka; Antikainen, Riitta; Tuomilehto, Jaakko (2007): Occupational, commuting, and leisure-time physical activity in relation to cardiovascular mortality among finnish subjects with hypertension. In: *American Journal of Hypertension* 20 (12), S. 1242–1250. DOI: 10.1016/j.amjhyper.2007.07.015.
- Huang, T.; Larsen, K. T.; Ried-Larsen, M.; Møller, N. C.; Andersen, L. B. (2014): The effects of physical activity and exercise on brain-derived neurotrophic factor in healthy humans: A review. In: *Scandinavian journal of medicine & science in sports* 24 (1), S. 1–10. DOI: 10.1111/sms.12069.
- Huang, Guoyuan; Shi, Xiangrong; Gibson, Cheryl A.; Huang, Sunny C.; Coudret, Nadine A.; Ehlman, Mary C. (2013): Controlled aerobic exercise training reduces resting blood pressure in sedentary older adults. In: *Blood Pressure* 22 (6), S. 386–394. DOI: 10.3109/08037051.2013.778003.
- Hulzebos, Erik; Dadema, Tessa; Takken, Tim (2013): Measurement of physical activity in patients with cystic fibrosis: a systematic review. In: *Expert Review of Respiratory Medicine* 7 (6), S. 647–653. DOI: 10.1586/17476348.2013.839248.
- Hunt, Sharon Ann; Abraham, William T.; Chin, Marshall H.; Feldman, Arthur M.; Francis, Gary S.; Ganiats, Theodore G. et al. (2005): ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. In: *Circulation* 112 (12), e154-235. DOI: 10.1161/CIRCULATIONAHA.105.167586.
- Hunter, Gary R.; Brock, David W.; Byrne, Nuala M.; Chandler-Laney, Paula C.; Del Corral, Pedro; Gower, Barbara A. (2010): Exercise training prevents regain of visceral fat for 1 year following weight loss. In: *Obesity (Silver Spring, Md.)* 18 (4), S. 690–695. DOI: 10.1038/oby.2009.316.
- Hurkmans (2009): Dynamic exercise programs (aerobic capacity and/or muscle strength training) in patients with rheumatoid arthritis. In: *Cochrane Database Syst Rev*, CD006853.
- Hurwitz, Eric L.; Morgenstern, Hal; Chiao, Chi (2005): Effects of recreational physical activity and back exercises on low back pain and psychological distress: findings from the UCLA Low Back Pain Study. In: *American Journal of Public Health* 95 (10), S. 1817–1824. DOI: 10.2105/AJPH.2004.052993.
- Hwang, Rita; Marwick, Thomas (2009): Efficacy of home-based exercise programmes for people with chronic heart failure: a meta-analysis. In: *European journal of cardiovascular prevention and rehabilitation : official journal of the European Society of Cardiology, Working Groups on Epidemiology & Prevention and Cardiac Rehabilitation and Exercise Physiology* 16 (5), S. 527–535. DOI: 10.1097/HJR.0b013e32832e097f.
- Ibrahim, Ezzeldin M.; Al-Homaidh, Abdelaziz (2011): Physical activity and survival after breast cancer diagnosis: meta-analysis of published studies. In: *Medical Oncology* 28 (3), S. 753–765. DOI: 10.1007/s12032-010-9536-x.
- Iepsen, Ulrik Winning; Jørgensen, Karsten Juhl; Ringbæk, Thomas; Hansen, Henrik; Skrubbeltrang, Conni; Lange, Peter (2015): A combination of resistance and endurance training increases leg muscle strength in COPD: An evidence-based recommendation based on systematic review with meta-analyses. In: *Chronic Respiratory Disease* 12 (2), S. 132–145. DOI: 10.1177/1479972315575318.
- Iepsen, Ulrik Winning; Jørgensen, Karsten Juhl; Ringbaek, Thomas; Hansen, Henrik; Skrubbeltrang, Conni; Lange, Peter (2015): A Systematic Review of Resistance Training Versus Endurance Training in COPD. In: *Journal of Cardiopulmonary Rehabilitation and Prevention* 35 (3), S. 163–172. DOI: 10.1097/HCR.000000000000105.
- Ihlebaek (2006): Prevalence of low back pain and sickness absence: a “borderline” study in Norway and Sweden. In: *Scand J Public Health* 34, S. 555.
- Ilanne-Parikka, Pirjo; Laaksonen, David E.; Eriksson, Johan G.; Lakka, Timo A.; Lindstr, Jaanaööm; Peltonen, Markku et al. (2010): Leisure-time physical activity and the metabolic syndrome in the Finnish diabetes prevention study. In: *Diabetes Care* 33 (7), S. 1610–1617. DOI: 10.2337/dc09-2155.
- Irvine, Casey; Taylor, Nicholas F. (2009): Progressive resistance exercise improves glycaemic control in people with type 2 diabetes mellitus: a systematic review. In: *Australian Journal of Physiotherapy* 55 (4), S. 237–246. DOI: 10.1016/S0004-9514(09)70003-0.
- Irwin, Melinda L.; Yasui, Yutaka; Ulrich, Cornelia M.; Bowen, Deborah; Rudolph, Rebecca E.; Schwartz, Robert S. et al. (2003): Effect of exercise on total and intra-abdominal body fat in postmenopausal women: a randomized controlled trial. In: *JAMA* 289 (3), S. 323–330. DOI: 10.1001/jama.289.3.323.

- Ismail, Hashbullah; McFarlane, James R.; Nojournian, A. Hadi; Dieberg, Gudrun; Smart, Neil A. (2013): Clinical outcomes and cardiovascular responses to different exercise training intensities in patients with heart failure: a systematic review and meta-analysis. In: *JACC. Heart failure* 1 (6), S. 514–522. DOI: 10.1016/j.jchf.2013.08.006.
- Iso-Markku, Paula; Waller, Katja; Kujala, Urho M.; Kaprio, Jaakko (2015): Physical activity and dementia: long-term follow-up study of adult twins. In: *Annals of Medicine* 47 (2), S. 81–87. DOI: 10.3109/07853890.2014.994675.
- IVY, JOHN L.; ZDERIC, T. W.E.D.; FOGT, DONOVAN L. (1999): 1 Prevention and Treatment of Non-Insulin-Dependent Diabetes Mellitus. In: *Exercise and Sport Sciences Reviews* 27, 1??36. DOI: 10.1249/00003677-199900270-00003.
- Iwamoto, J.; Takeda, T.; Ichimura, S. (2001): Effect of exercise training and detraining on bone mineral density in postmenopausal women with osteoporosis. In: *Journal of Orthopaedic Science* 6 (2), S. 128–132. DOI: 10.1007/s007760100059.
- Iwata, Masaaki; Ota, Kristie T.; Duman, Ronald S. (2013): The inflammasome: pathways linking psychological stress, depression, and systemic illnesses. In: *Brain, Behavior, and Immunity* 31, S. 105–114. DOI: 10.1016/j.bbi.2012.12.008.
- Jakubowska, Joanna; Bohdanowicz-Pawlak, Anna; Milewicz, Andrzej; Szymczak, Jadwiga; Bednarek-Tupikowska, Grazyna; Demissie, Marek (2008): Plasma cytokines in obese women with polycystic ovary syndrome, before and after metformin treatment. In: *Gynecological Endocrinology* 24 (7), S. 378–384. DOI: 10.1080/09513590802128968.
- Janiszewski, P. M.; Kuk, J. L.; Ross, R. (2008): Is the reduction of lower-body subcutaneous adipose tissue associated with elevations in risk factors for diabetes and cardiovascular disease? In: *Diabetologia* 51 (8), S. 1475–1482. DOI: 10.1007/s00125-008-1058-0.
- Janiszewski, Peter M.; Ross, Robert (2007): Physical activity in the treatment of obesity: beyond body weight reduction. In: *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme* 32 (3), S. 512–522. DOI: 10.1139/H07-018.
- Janssen, Ian; Fortier, Anne; Hudson, Robert; Ross, Robert (2002): Effects of an energy-restrictive diet with or without exercise on abdominal fat, intermuscular fat, and metabolic risk factors in obese women. In: *Diabetes Care* 25 (3), S. 431–438. DOI: 10.2337/diacare.25.3.431.
- Jensen, Ole Kudsk; Nielsen, Claus Vinther; Stengaard-Pedersen, Kristian (2010): Low back pain may be caused by disturbed pain regulation: a cross-sectional study in low back pain patients using tender point examination. In: *European journal of pain (London, England)* 14 (5), S. 514–522. DOI: 10.1016/j.ejpain.2009.09.002.
- Jewitt, D. E.; Reid, D.; Thomas, M.; Mercer, C. J.; Valori, C.; Shillingford, J. P. (1969): FREE NORADRENALINE AND ADRENALINE EXCRETION IN RELATION TO THE DEVELOPMENT OF CARDIAC ARRHYTHMIAS AND HEART-FAILURE IN PATIENTS WITH ACUTE MYOCARDIAL INFARCTION. In: *The Lancet* 293 (7596), S. 635–641. DOI: 10.1016/S0140-6736(69)92009-1.
- Johns, David J.; Hartmann-Boyce, Jamie; Jebb, Susan A.; Aveyard, Paul (2014): Diet or exercise interventions vs combined behavioral weight management programs: a systematic review and meta-analysis of direct comparisons. In: *Journal of the Academy of Nutrition and Dietetics* 114 (10), S. 1557–1568. DOI: 10.1016/j.jand.2014.07.005.
- Johnstone, M. T.; Creager, S. J.; Scales, K. M.; Cusco, J. A.; Lee, B. K.; Creager, M. A. (1993): Impaired endothelium-dependent vasodilation in patients with insulin-dependent diabetes mellitus. In: *Circulation* 88 (6), S. 2510–2516. DOI: 10.1161/01.CIR.88.6.2510.
- Jolliffe (2000): Exercise-based rehabilitation for coronary heart disease. In: *Cochrane Database Syst Rev* 4, CD001800.
- Jonason (1987): Prediction of the effect of training on the walking tolerance in patients with intermittent claudication. In: *Scand J Rehabil Med* 19, S. 47.
- Jong, Zuzana de; Munneke, Marten; Zwinderman, Aeilko H.; Kroon, Herman M.; Jansen, Annemarie; Runday, Karel H. et al. (2003): Is a long-term high-intensity exercise program effective and safe in patients with rheumatoid arthritis? Results of a randomized controlled trial. In: *Arthritis and rheumatism* 48 (9), S. 2415–2424. DOI: 10.1002/art.11216.
- Jong, Z. de; Munneke, M.; Zwinderman, A. H.; Kroon, H. M.; Runday, K. H.; Lems, W. F. et al. (2004): Long term high intensity exercise and damage of small joints in rheumatoid arthritis. In: *Annals of the Rheumatic Diseases* 63 (11), S. 1399–1405. DOI: 10.1136/ard.2003.015826.
- Jong, Zuzana de; Munneke, Marten; Lems, Willem F.; Zwinderman, Aeilko H.; Kroon, Herman M.; Pauwels, Ernest K. J. et al. (2004): Slowing of bone loss in patients with rheumatoid arthritis by long-term high-intensity exercise: results of a randomized, controlled trial. In: *Arthritis and rheumatism* 50 (4), S. 1066–1076. DOI: 10.1002/art.20117.
- Josefsson, T.; Lindwall, M.; Archer, T. (2014): Physical exercise intervention in depressive disorders: meta-analysis and systematic review. In: *Scandinavian journal of medicine & science in sports* 24 (2), S. 259–272. DOI: 10.1111/sms.12050.
- Joslin (1959): The treatment of diabetes mellitus.
- Juhl, C.; Christensen, R.; Roos, E. M.; Zhang, W.; Lund, H. (2014): Impact of exercise type and dose on pain and disability in knee osteoarthritis: a systematic review and meta-regression analysis of randomized controlled trials. In: *Arthritis & rheumatology (Hoboken, N.J.)* 66 (3), S. 622–636. DOI: 10.1002/art.38290.

- Kamper, Steven J.; Apeldoorn, A. T.; Chiarotto, A.; Smeets, R. J. E. M.; Ostelo, R. W. J. G.; Guzman, J.; van Tulder, M. W. (2015): Multidisciplinary biopsychosocial rehabilitation for chronic low back pain: Cochrane systematic review and meta-analysis. In: *BMJ* 350 (feb18 5), h444. DOI: 10.1136/bmj.h444.
- Kanis, J. A. (1993): The incidence of hip fracture in Europe. In: *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 3 Suppl 1 (S1), S. 10–15. DOI: 10.1007/BF01621853.
- Kannel, William B. (1979): Diabetes and Cardiovascular Disease. In: *JAMA* 241 (19), S. 2035. DOI: 10.1001/jama.1979.03290450033020.
- Karila, C.; Ravilly, S.; Gauthier, R.; Tardif, C.; Neveu, H.; Maire, J. et al. (2010): Activité physique et réentraînement à l'effort du patient atteint de mucoviscidose. In: *Revue des Maladies Respiratoires* 27 (4), S. 301–313. DOI: 10.1016/j.rmr.2010.02.012.
- Karstoft, Kristian; Christensen, Camilla S.; Pedersen, Bente K.; Solomon, Thomas P. J. (2014): The acute effects of interval- Vs continuous-walking exercise on glycemic control in subjects with type 2 diabetes: a crossover, controlled study. In: *The Journal of Clinical Endocrinology & Metabolism* 99 (9), S. 3334–3342. DOI: 10.1210/jc.2014-1837.
- Karstoft, Kristian; Winding, Kamilla; Knudsen, Sine H.; Nielsen, Jens S.; Thomsen, Carsten; Pedersen, Bente K.; Solomon, Thomas P. J. (2013): The effects of free-living interval-walking training on glycemic control, body composition, and physical fitness in type 2 diabetic patients: a randomized, controlled trial. In: *Diabetes Care* 36 (2), S. 228–236. DOI: 10.2337/dc12-0658.
- Kayman, S.; Bruvold, W.; Stern, J. S. (1990): Maintenance and relapse after weight loss in women: behavioral aspects. In: *The American Journal of Clinical Nutrition* 52 (5), S. 800–807. DOI: 10.1093/ajcn/52.5.800.
- Kearney (2015): Exercise and postprandial glycemic control in type 2 diabetes. In: *Curr Diabetes Rev.*
- Kelemen, Michael H. (1990): Exercise Training Combined With Antihypertensive Drug Therapy. In: *JAMA* 263 (20), S. 2766. DOI: 10.1001/jama.1990.03440200070025.
- Kelestimur, Fahrettin; Unluhizarci, Kürşad; Baybuga, Hür; Atmaca, Hulusi; Bayram, Fahri; Sahin, Yilmaz (2006): Prevalence of polycystic ovarian changes and polycystic ovary syndrome in premenopausal women with treated type 2 diabetes mellitus. In: *Fertility and Sterility* 86 (2), S. 405–410. DOI: 10.1016/j.fertnstert.2006.01.019.
- Kelley, G. A.; Kelley, K. S. (2000): Progressive resistance exercise and resting blood pressure : A meta-analysis of randomized controlled trials. In: *Hypertension* 35 (3), S. 838–843. DOI: 10.1161/01.HYP.35.3.838.
- Kelley, G. A.; Kelley, K. S. (2007): Effects of aerobic exercise on lipids and lipoproteins in adults with type 2 diabetes: a meta-analysis of randomized-controlled trials. In: *Public Health* 121 (9), S. 643–655. DOI: 10.1016/j.puhe.2007.02.014.
- Kelley, George A.; Kelley, Kristi S. (2010): Isometric handgrip exercise and resting blood pressure: a meta-analysis of randomized controlled trials. In: *Journal of Hypertension* 28 (3), S. 411–418. DOI: 10.1097/HJH.0b013e3283357d16.
- Kelley, G. A.; Kelley, K. A.; Tran, Z. V. (2001): Aerobic exercise and resting blood pressure: a meta-analytic review of randomized, controlled trials. In: *Preventive Cardiology* 4 (2), S. 73–80. DOI: 10.1111/j.1520-037X.2001.00529.x.
- Kelley, George A.; Kelley, Kristi S.; Hootman, Jennifer M.; Jones, Dina L. (2011): Effects of community-deliverable exercise on pain and physical function in adults with arthritis and other rheumatic diseases: a meta-analysis. In: *Arthritis Care & Research* 63 (1), S. 79–93. DOI: 10.1002/acr.20347.
- Kemoun, Gilles; Thibaud, Marie; Roumagne, Nicolas; Carette, Pierre; Albinet, Cédric; Toussaint, Lucette et al. (2010): Effects of a physical training programme on cognitive function and walking efficiency in elderly persons with dementia. In: *Dementia and Geriatric Cognitive Disorders* 29 (2), S. 109–114. DOI: 10.1159/000272435.
- Kemper, H.C.G.; Twisk, J.W.R.; van Mechelen, W.; Post, G.B.; Roos, J.C.; Lips, P. (2000): A fifteen-year longitudinal study in young adults on the relation of physical activity and fitness with the development of the bone mass: the Amsterdam Growth and Health Longitudinal Study. In: *Bone* 27 (6), S. 847–853. DOI: 10.1016/S8756-3282(00)00397-5.
- Kenfield, Stacey A.; Stampfer, Meir J.; Giovannucci, Edward; Chan, June M. (2011): Physical activity and survival after prostate cancer diagnosis in the health professionals follow-up study. In: *Journal of Clinical Oncology* 29 (6), S. 726–732. DOI: 10.1200/JCO.2010.31.5226.
- Kennedy, Amy; Nirantharakumar, Krishnarajah; Chimen, Myriam; Pang, Terence T.; Hemming, Karla; Andrews, Rob C.; Narendran, Parth (2013): Does exercise improve glycaemic control in type 1 diabetes? A systematic review and meta-analysis. In: *PLoS ONE* 8 (3), e58861. DOI: 10.1371/journal.pone.0058861.
- Kent-Braun, J. A.; Sharma, K. R.; Weiner, M. W.; Miller, R. G. (1994): Effects of exercise on muscle activation and metabolism in multiple sclerosis. In: *Muscle & Nerve* 17 (10), S. 1162–1169. DOI: 10.1002/mus.880171006.
- Keogh, Justin W. L.; MacLeod, Roderick D. (2012): Body composition, physical fitness, functional performance, quality of life, and fatigue benefits of exercise for prostate cancer patients: a systematic review. In: *Journal of Pain and Symptom Management* 43 (1), S. 96–110. DOI: 10.1016/j.jpainsymman.2011.03.006.
- Khan (2007): Multidisciplinary rehabilitation for adults with multiple sclerosis. In: *Cochrane Database Syst Rev*, CD006036.



- Kiilavuori, K.; Sovijärvi, A.; Näveri, H.; Ikonen, T.; Leinonen, H. (1996): Effect of physical training on exercise capacity and gas exchange in patients with chronic heart failure. In: *Chest* 110 (4), S. 985–991. DOI: 10.1378/chest.110.4.985.
- Kiilavuori, K.; TOIVONEN, L.; Näveri, H.; Leinonen, H. (1995): Reversal of autonomic derangements by physical training in chronic heart failure assessed by heart rate variability. In: *European Heart Journal* 16 (4), S. 490–495. DOI: 10.1093/oxfordjournals.eurheartj.a060941.
- Kim, Jae-Min; Stewart, Robert; Bae, Kyung-Yeol; Kim, Sung-Wan; Yang, Su-Jin; Park, Kee-Hyung et al. (2011): Role of BDNF val66met polymorphism on the association between physical activity and incident dementia. In: *Neurobiology of Aging* 32 (3), 551.e5-12. DOI: 10.1016/j.neurobiolaging.2010.01.018.
- King, Abby C. (1989): Diet vs Exercise in Weight Maintenance. In: *Archives of Internal Medicine* 149 (12), S. 2741. DOI: 10.1001/archinte.1989.00390120085017.
- King, Marjorie L.; Guarracini, Mary; Lennihan, Laura; Freeman, Deborah; Gagas, Barbara; Boston, Andrew et al. (1989): Adaptive Exercise Testing for Patients with Hemiparesis. In: *Journal of Cardiopulmonary Rehabilitation* 9 (6), S. 237–242. DOI: 10.1097/00008483-198906000-00002.
- Kirk, Alison; Mutrie, Nanette; MacIntyre, Paul; Fisher, Miles (2003): Increasing physical activity in people with type 2 diabetes. In: *Diabetes Care* 26 (4), S. 1186–1192. DOI: 10.2337/diacare.26.4.1186.
- Kjølhede, T.; Vissing, K.; Dalgas, U. (2012): Multiple sclerosis and progressive resistance training: a systematic review. In: *Multiple Sclerosis Journal* 18 (9), S. 1215–1228. DOI: 10.1177/1352458512437418.
- Klijn, Peter H. C.; Oudshoorn, Annemarie; van der Ent, Cornelis K.; van der Net, Janjaap; Kimpen, Jan L.; Helden, Paul J. M. (2004): Effects of anaerobic training in children with cystic fibrosis: a randomized controlled study. In: *Chest* 125 (4), S. 1299–1305. DOI: 10.1378/chest.125.4.1299.
- Knopp, R. H. (1999): Drug treatment of lipid disorders. In: *The New England journal of medicine* 341 (7), S. 498–511. DOI: 10.1056/NEJM199908123410707.
- Knowler, William C.; Barrett-Connor, Elizabeth; Fowler, Sarah E.; Hamman, Richard F.; Lachin, John M.; Walker, Elizabeth A.; Nathan, David M. (2002): Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. In: *New England Journal of Medicine* 346 (6), S. 393–403. DOI: 10.1056/NEJMoa012512.
- Kodama, Satoru; Tanaka, Shiro; Saito, Kazumi; Shu, Miao; Sone, Yasuko; Onitake, Fumiko et al. (2007): Effect of aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a meta-analysis. In: *Archives of Internal Medicine* 167 (10), S. 999–1008. DOI: 10.1001/archinte.167.10.999.
- Kohl, H. W.; Gordon, N. F.; Villegas, J. A.; Blair, S. N. (1992): Cardiorespiratory fitness, glycemic status, and mortality risk in men. In: *Diabetes Care* 15 (2), S. 184–192. DOI: 10.2337/diacare.15.2.184.
- Koivisto, V. A.; Felig, P. (1978): Effects of leg exercise on insulin absorption in diabetic patients. In: *The New England journal of medicine* 298 (2), S. 79–83. DOI: 10.1056/NEJM197801122980205.
- Konerth, Megan; Childers, Jennifer (2013): Exercise: a possible adjunct therapy to alleviate early Parkinson disease. In: *Journal of the American Academy of Physician Assistants* 26 (4), S. 30–33. DOI: 10.1097/01720610-201304000-00007.
- Kongsgaard, M.; Backer, V.; Jørgensen, K.; Kjaer, M.; Beyer, N. (2004): Heavy resistance training increases muscle size, strength and physical function in elderly male COPD-patients--a pilot study. In: *Respiratory Medicine* 98 (10), S. 1000–1007. DOI: 10.1016/j.rmed.2004.03.003.
- Krørup, Lars-Henrik; Truelsen, Thomas; Pedersen, Anders; Lerke, Hanne; Lindahl, Marianne; Hansen, Lotte et al. (2007): Level of physical activity in the week preceding an ischemic stroke. In: *Cerebrovascular Diseases* 24 (2-3), S. 296–300. DOI: 10.1159/000105683.
- Krørup, L-H; Truelsen, T.; Gluud, C.; Andersen, G.; Zeng, X.; Kõrv, J. et al. (2008): Prestroke physical activity is associated with severity and long-term outcome from first-ever stroke. In: *Neurology* 71 (17), S. 1313–1318. DOI: 10.1212/01.wnl.0000327667.48013.9f.
- Kraus, William E.; Houmard, Joseph A.; Duscha, Brian D.; Knetzger, Kenneth J.; Wharton, Michelle B.; McCartney, Jennifer S. et al. (2002): Effects of the amount and intensity of exercise on plasma lipoproteins. In: *New England Journal of Medicine* 347 (19), S. 1483–1492. DOI: 10.1056/NEJMoa020194.
- Krolewski, Andrzej S.; Kosinski, Edward J.; Warram, James H.; Stevens Leland, O.; Busick, Edward J.; Cader Asmal, A. et al. (1987): Magnitude and determinants of coronary artery disease in juvenile-onset, insulin-dependent diabetes mellitus. In: *The American Journal of Cardiology* 59 (8), S. 750–755. DOI: 10.1016/0002-9149(87)91086-1.
- Krølner, B.; Toft, B. (1983): Vertebral bone loss: an unheeded side effect of therapeutic bed rest. In: *Clinical science (London, England : 1979)* 64 (5), S. 537–540. DOI: 10.1042/cs0640537.
- Krotkiewski, M.; Lönnroth, P.; Mandroukas, K.; Wroblewski, Z.; Rebuffé-Scrive, M.; Holm, G. et al. (1985): The effects of physical training on insulin secretion and effectiveness and on glucose metabolism in obesity and type 2 (non-insulin-dependent) diabetes mellitus. In: *Diabetologia* 28 (12), S. 881–890. DOI: 10.1007/BF00703130.

- Kuk, Jennifer L.; Katzmarzyk, Peter T.; Nichaman, Milton Z.; Church, Timothy S.; Blair, Steven N.; Ross, Robert (2006): Visceral fat is an independent predictor of all-cause mortality in men. In: *Obesity* 14 (2), S. 336–341. DOI: 10.1038/oby.2006.43.
- Laaksonen, D. E.; Atalay, M.; Niskanen, L. K.; Mustonen, J.; Sen, C. K.; Lakka, T. A.; Uusitupa, M. I. (2000): Aerobic exercise and the lipid profile in type 1 diabetic men: a randomized controlled trial. In: *Medicine & Science in Sports & Exercise* 32 (9), S. 1541–1548. DOI: 10.1097/00005768-200009000-00003.
- Lacasse (2002): Pulmonary rehabilitation for chronic obstructive pulmonary disease. In: *Cochrane Database Syst Rev* 3, CD003793.
- Lacasse (2007): Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. A Cochrane systematic review. In: *Eura Medicophys* 43, S. 475.
- Lacasse, Yves; Wong, Eric; Guyatt, Gordon H.; King, Derek; Cook, Deborah J.; Goldstein, Roger S. (1996): Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. In: *The Lancet* 348 (9035), S. 1115–1119. DOI: 10.1016/S0140-6736(96)04201-8.
- Landin (1977): Muscle metabolism during exercise in hemiparetic patients. In: *Clin Sci Mol Med* 53, S. 257.
- Lane (2014): Exercise for intermittent claudication. In: *Cochrane Database Syst Rev* 7, CD000990.
- Langbein, W. Edwin; Collins, Eileen G.; Orebaugh, Cynthia; Maloney, Christine; Williams, Karla J.; Littooy, Fred N.; Edwards, Lonnie C. (2002): Increasing exercise tolerance of persons limited by claudication pain using polestriding. In: *Journal of Vascular Surgery* 35 (5), S. 887–893. DOI: 10.1067/mva.2002.123756.
- Lange, Angela K.; Vanwanseele, Benedicte; Fiatarone singh, Maria A. (2008): Strength training for treatment of osteoarthritis of the knee: a systematic review. In: *Arthritis and rheumatism* 59 (10), S. 1488–1494. DOI: 10.1002/art.24118.
- Lannefors (2004): Physiotherapy in infants and young children with cystic fibrosis: current practice and future developments. In: *J R Soc Med* 97 (Suppl. 44), S. 8.
- LARRYDURSTINE, J.; HASKELL, WILLIAM L. (1994): Effects of Exercise Training on Plasma Lipids and Lipoproteins. In: *Exercise and Sport Sciences Reviews* 22 (1), S. 477–522. DOI: 10.1249/00003677-199401000-00017.
- Larson, Eric B.; Wang, Li; Bowen, James D.; McCormick, Wayne C.; Teri, Linda; Crane, Paul; Kukull, Walter (2006): Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. In: *Annals of Internal Medicine* 144 (2), S. 73–81. DOI: 10.7326/0003-4819-144-2-200601170-00004.
- Laske, Christoph; Banschbach, Sabine; Stransky, Elke; Bosch, Sabine; Straten, Guido; Machann, Jürgen et al. (2010): Exercise-induced normalization of decreased BDNF serum concentration in elderly women with remitted major depression. In: *The International Journal of Neuropsychopharmacology* 13 (5), S. 595–602. DOI: 10.1017/S1461145709991234.
- Latimer-Cheung, Amy E.; Pilutti, Lara A.; Hicks, Audrey L.; Martin Ginis, Kathleen A.; Fenuta, Alyssa M.; MacKibbin, K. Ann; Motl, Robert W. (2013): Effects of exercise training on fitness, mobility, fatigue, and health-related quality of life among adults with multiple sclerosis: a systematic review to inform guideline development. In: *Arch Phys Med Rehabil* 94 (9), 1800-1828.e3. DOI: 10.1016/j.apmr.2013.04.020.
- Lau, Lonneke M. L. de; Breteler, Monique M. B. (2006): Epidemiology of Parkinson's disease. In: *The Lancet Neurology* 5 (6), S. 525–535. DOI: 10.1016/S1474-4422(06)70471-9.
- Laurin, D.; Verreault, R.; Lindsay, J.; MacPherson, K.; Rockwood, K. (2001): Physical activity and risk of cognitive impairment and dementia in elderly persons. In: *Archives of Neurology* 58 (3), S. 498–504. DOI: 10.1001/archneur.58.3.498.
- Laursen, Thomas Munk; Nordentoft, Merete (2011): Heart disease treatment and mortality in schizophrenia and bipolar disorder - changes in the Danish population between 1994 and 2006. In: *Journal of Psychiatric Research* 45 (1), S. 29–35. DOI: 10.1016/j.jpsychires.2010.04.027.
- Lautenschlager, Nicola T.; Cox, Kay L.; Flicker, Leon; Foster, Jonathan K.; van Bockxmeer, Frank M.; Xiao, Jianguo et al. (2008): Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. In: *JAMA* 300 (9), S. 1027–1037. DOI: 10.1001/jama.300.9.1027.
- Law, M. R.; Wald, N. J.; Meade, T. W. (1991): Strategies for prevention of osteoporosis and hip fracture. In: *BMJ* 303 (6800), S. 453–459. DOI: 10.1136/bmj.303.6800.453.
- Lee, Chong Do; Folsom, Aaron R.; Blair, Steven N. (2003): Physical activity and stroke risk: a meta-analysis. In: *Stroke* 34 (10), S. 2475–2481. DOI: 10.1161/01.STR.0000091843.02517.9D.
- Lee, SoJung; Kuk, Jennifer L.; Davidson, Lance E.; Hudson, Robert; Kilpatrick, Katherine; Graham, Terry E.; Ross, Robert (2005): Exercise without weight loss is an effective strategy for obesity reduction in obese individuals with and without Type 2 diabetes. In: *Journal of applied physiology (Bethesda, Md. : 1985)* 99 (3), S. 1220–1225. DOI: 10.1152/jappphysiol.00053.2005.
- Leermakers, Elizabeth A.; Perri, Michael G.; Shigaki, Cheryl L.; Fuller, Pamela R. (1999): Effects of exercise-focused versus weight-focused maintenance programs on the management of obesity. In: *Addictive Behaviors* 24 (2), S. 219–227. DOI: 10.1016/S0306-4603(98)00090-2.

- Lehmann, R.; Engler, H.; Honegger, R.; Riesen, W.; Spinass, G. A. (2001): Alterations of lipolytic enzymes and high-density lipoprotein subfractions induced by physical activity in type 2 diabetes mellitus. In: *European Journal of Clinical Investigation* 31 (1), S. 37–44. DOI: 10.1046/j.1365-2362.2001.00752.x.
- Lehmann, R.; Kaplan, V.; Bingisser, R.; Bloch, K. E.; Spinass, G. A. (1997): Impact of physical activity on cardiovascular risk factors in IDDM. In: *Diabetes Care* 20 (10), S. 1603–1611. DOI: 10.2337/diacare.20.10.1603.
- Lehmann, R.; Vokac, A.; Niedermann, K.; Agosti, K.; Spinass, G. A. (1995): Loss of abdominal fat and improvement of the cardiovascular risk profile by regular moderate exercise training in patients with NIDDM. In: *Diabetologia* 38 (11), S. 1313–1319. DOI: 10.1007/BF00401764.
- LeMaitre, John P.; Harris, Stuart; Fox, Keith A. A.; Denvir, Martin (2004): Change in circulating cytokines after 2 forms of exercise training in chronic stable heart failure. In: *American Heart Journal* 147 (1), S. 100–105. DOI: 10.1016/j.ahj.2003.07.001.
- Lemmey, Andrew B.; Marcora, Samuele M.; Chester, Kathryn; Wilson, Sally; Casanova, Francesco; Maddison, Peter J. (2009): Effects of high-intensity resistance training in patients with rheumatoid arthritis: a randomized controlled trial. In: *Arthritis and rheumatism* 61 (12), S. 1726–1734. DOI: 10.1002/art.24891.
- Leon, Arthur S. (1991): Effects of Exercise Conditioning on Physiologic Precursors of Coronary Heart Disease. In: *Journal of Cardiopulmonary Rehabilitation* 11 (1), S. 46–57. DOI: 10.1097/00008483-199101000-00008.
- Leon (1999): Exercise and the heart in health and disease.
- Leon, A. S.; Sanchez, O. A. (2001): Response of blood lipids to exercise training alone or combined with dietary intervention. In: *Medicine & Science in Sports & Exercise* 33 (6 Suppl), S502–15; discussion S528–9. DOI: 10.1097/00005768-200106001-00021.
- Levy, W. C.; Cerqueira, M. D.; Abrass, I. B.; Schwartz, R. S.; Stratton, J. R. (1993): Endurance exercise training augments diastolic filling at rest and during exercise in healthy young and older men. In: *Circulation* 88 (1), S. 116–126. DOI: 10.1161/01.CIR.88.1.116.
- LeWitt, P. A.; Bharucha, A.; Chitrit, I.; Takis, C.; Patil, S.; Schork, M. A.; Pichurko, B. (1994): Perceived exertion and muscle efficiency in Parkinson's disease: L-DOPA effects. In: *Clinical Neuropharmacology* 17 (5), S. 454–459. DOI: 10.1097/00002826-199410000-00008.
- Li (2014): Diabetes mellitus and dementia - a systematic review and meta-analysis. In: *Eur Rev Med Pharmacol Sci* 18, S. 1778.
- Li, Y. P.; Reid, M. B. (2001): Effect of tumor necrosis factor- $\alpha$  on skeletal muscle metabolism. In: *Current Opinion in Rheumatology* 13 (6), S. 483–487. DOI: 10.1097/00002281-200111000-00005.
- Liang, Hui; Zhang, Hua; Ji, Haiyan; Wang, Chunmei (2015): Effects of home-based exercise intervention on health-related quality of life for patients with ankylosing spondylitis: a meta-analysis. In: *Clinical Rheumatology* 34 (10), S. 1737–1744. DOI: 10.1007/s10067-015-2913-2.
- Lindegård, B.; Hansen, T.; Hvid, T.; van Hall, G.; Plomgaard, P.; Ditlevsen, S. et al. (2008): The effect of strength and endurance training on insulin sensitivity and fat distribution in human immunodeficiency virus-infected patients with lipodystrophy. In: *The Journal of clinical endocrinology and metabolism* 93 (10), S. 3860–3869. DOI: 10.1210/jc.2007-2733.
- Lindenstrøm, E.; Boysen, G.; Nyboe, J. (1993): Lifestyle factors and risk of cerebrovascular disease in women. The Copenhagen City Heart Study. In: *Stroke* 24 (10), S. 1468–1472. DOI: 10.1161/01.STR.24.10.1468.
- Lindhardsen (2011): [Excess mortality from cardiovascular disease in patients with rheumatoid arthritis]. In: *Ugeskr Laeger* 173, S. 343.
- Lindström, Jaana; Eriksson, Johan G.; Valle, Timo T.; Aunola, Sirkka; Cepaitis, Zygimantas; Hakumäki, Martti et al. (2003): Prevention of diabetes mellitus in subjects with impaired glucose tolerance in the Finnish Diabetes Prevention Study: results from a randomized clinical trial. In: *Journal of the American Society of Nephrology : JASN* 14 (7 Suppl 2), S108–13. DOI: 10.1097/01.ASN.0000070157.96264.13.
- Lindström, Jaana; Louheranta, Anne; Mannelin, Marjo; Rastas, Merja; Salminen, Virpi; Eriksson, Johan et al. (2003): The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity. In: *Diabetes Care* 26 (12), S. 3230–3236. DOI: 10.2337/diacare.26.12.3230.
- Little, Jonathan P.; Gillen, Jenna B.; Percival, Michael E.; Safdar, Adeel; Tarnopolsky, Mark A.; Punthakee, Zubin et al. (2011): Low-volume high-intensity interval training reduces hyperglycemia and increases muscle mitochondrial capacity in patients with type 2 diabetes. In: *Journal of Applied Physiology* 111 (6), S. 1554–1560. DOI: 10.1152/jappphysiol.00921.2011.
- Lloyd-Williams (2002): Exercise training and heart failure: a systematic review of current evidence. In: *Br J Gen Pract* 52, S. 47.
- Loge, J. H.; Abrahamsen, A. F.; Ekeberg, O.; Kaasa, S. (1999): Hodgkin's disease survivors more fatigued than the general population. In: *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 17 (1), S. 253–261. DOI: 10.1200/JCO.1999.17.1.253.
- Lokey, E. A.; Tran, Z. V. (1989): Effects of exercise training on serum lipid and lipoprotein concentrations in women: a meta-analysis. In: *International Journal of Sports Medicine* 10 (6), S. 424–429. DOI: 10.1055/s-2007-1024937.

- Long, Audrey; Donelson, Ron; Fung, Tak (2004): Does it matter which exercise? A randomized control trial of exercise for low back pain. In: *Spine* 29 (23), S. 2593–2602. DOI: 10.1097/01.brs.0000146464.23007.2a.
- LORENZ, LANDON B.; WILD, ROBERT A. (2007): Polycystic ovarian syndrome: an evidence-based approach to evaluation and management of diabetes and cardiovascular risks for today's clinician. In: *Clinical Obstetrics and Gynecology* 50 (1), S. 226–243. DOI: 10.1097/GRF.0b013e31802f5197.
- Love, Caitlin; Sabiston, Catherine M. (2011): Exploring the links between physical activity and posttraumatic growth in young adult cancer survivors. In: *Psycho-Oncology* 20 (3), S. 278–286. DOI: 10.1002/pon.1733.
- Lukoff, D.; Wallace, C. J.; Liberman, R. P.; Burke, K. (1986): A holistic program for chronic schizophrenic patients. In: *Schizophrenia Bulletin* 12 (2), S. 274–282. DOI: 10.1093/schbul/12.2.274.
- Lunde, Ottar; Tanbo, Tom (2007): Polycystic ovary syndrome: a follow-up study on diabetes mellitus, cardiovascular disease and malignancy 15–25 years after ovarian wedge resection. In: *Gynecological endocrinology : the official journal of the International Society of Gynecological Endocrinology* 23 (12), S. 704–709. DOI: 10.1080/09513590701705189.
- Lundgren, F.; Dahllöf, A. G.; Scherstén, T.; Bylund-Fellenius, A. C. (1989): Muscle enzyme adaptation in patients with peripheral arterial insufficiency: spontaneous adaptation, effect of different treatments and consequences on walking performance. In: *Clinical science (London, England : 1979)* 77 (5), S. 485–493. DOI: 10.1042/cs0770485.
- Lyngberg (1988): The effect of physical training on patients with rheumatoid arthritis: changes in disease activity, muscle strength and aerobic capacity. A clinically controlled minimized cross-over study. In: *Clin Exp Rheumatol* 6, S. 253.
- Lyngberg, Kirsten K.; Harreby, Mette; Bentzen, Helen; Frost, Bodil; Danneskiold-Samsøe, Bente (1994): Elderly rheumatoid arthritis patients on steroid treatment tolerate physical training without an increase in disease activity. In: *Archives of Physical Medicine and Rehabilitation* 75 (11), S. 1189–1195. DOI: 10.1016/0003-9993(94)90003-5.
- Lytle, Mary Ellen; Vander Bilt, Joni; Pandav, Rajesh S.; Dodge, Hiroko H.; Ganguli, Mary (2004): Exercise level and cognitive decline: the MoVIES project. In: *Alzheimer Disease & Associated Disorders* 18 (2), S. 57–64. DOI: 10.1097/01.wad.0000126614.87955.79.
- MacLeod, S. F.; Terada, T.; Chahal, B. S.; Boulé, N. G. (2013): Exercise lowers postprandial glucose but not fasting glucose in type 2 diabetes: a meta-analysis of studies using continuous glucose monitoring. In: *Diabetes/Metabolism Research and Reviews* 29 (8), S. 593–603. DOI: 10.1002/dmrr.2461.
- Madureira, Melisa M.; Bonfá, Eloisa; Takayama, Liliam; Pereira, Rosa M. R. (2010): A 12-month randomized controlled trial of balance training in elderly women with osteoporosis: improvement of quality of life. In: *Maturitas* 66 (2), S. 206–211. DOI: 10.1016/j.maturitas.2010.03.009.
- Madureira, M. M.; Takayama, L.; Gallinaro, A. L.; Caparbo, V. F.; Costa, R. A.; Pereira, R. M. R. (2007): Balance training program is highly effective in improving functional status and reducing the risk of falls in elderly women with osteoporosis: a randomized controlled trial. In: *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* 18 (4), S. 419–425. DOI: 10.1007/s00198-006-0252-5.
- Mak, Winifred; Dokras, Anuja (2009): Polycystic ovarian syndrome and the risk of cardiovascular disease and thrombosis. In: *Seminars in Thrombosis and Hemostasis* 35 (7), S. 613–620. DOI: 10.1055/s-0029-1242715.
- Mäkimattila, S.; Mäntysaari, M.; Groop, P. H.; Summanen, P.; Virkamäki, A.; Schlenzka, A. et al. (1997): Hyperreactivity to nitrovasodilators in forearm vasculature is related to autonomic dysfunction in insulin-dependent diabetes mellitus. In: *Circulation* 95 (3), S. 618–625. DOI: 10.1161/01.CIR.95.3.618.
- Mäkimattila, S.; Virkamäki, A.; Groop, P. H.; Cockcroft, J.; Utriainen, T.; Fagerudd, J.; Yki-Järvinen, H. (1996): Chronic hyperglycemia impairs endothelial function and insulin sensitivity via different mechanisms in insulin-dependent diabetes mellitus. In: *Circulation* 94 (6), S. 1276–1282. DOI: 10.1161/01.CIR.94.6.1276.
- Mälkiä, E.; Impivaara, O. (1998): Intensity of physical activity and respiratory function in subjects with and without bronchial asthma. In: *Scandinavian journal of medicine & science in sports* 8 (1), S. 27–32. DOI: 10.1111/j.1600-0838.1998.tb00225.x.
- Malmivaara, Antti; Häkkinen, Unto; Aro, Timo; Heinrichs, Maj-Len; Koskeniemi, Liisa; Kuosma, Eeva et al. (1995): The Treatment of Acute Low Back Pain — Bed Rest, Exercises, or Ordinary Activity? In: *New England Journal of Medicine* 332 (6), S. 351–355. DOI: 10.1056/NEJM199502093320602.
- Mandroukas, K.; Krotkiewski, M.; Hedberg, M.; Wroblewski, Z.; Björntorp, P.; Grimby, G. (1984): Physical training in obese women. Effects of muscle morphology, biochemistry and function. In: *European Journal of Applied Physiology and Occupational Physiology* 52 (4), S. 355–361. DOI: 10.1007/BF00943363.
- Manji, Hussein K.; Moore, Gregory J.; Chen, Guang (2000): Clinical and preclinical evidence for the neurotrophic effects of mood stabilizers: implications for the pathophysiology and treatment of manic–depressive illness. In: *Biological Psychiatry* 48 (8), S. 740–754. DOI: 10.1016/S0006-3223(00)00979-3.
- Mann, Steven; Beedie, Christopher; Jimenez, Alfonso (2014): Differential effects of aerobic exercise, resistance training and combined exercise modalities on cholesterol and the lipid profile: review, synthesis and recommendations. In: *Sports medicine (Auckland, N.Z.)* 44 (2), S. 211–221. DOI: 10.1007/s40279-013-0110-5.

- Mannarino, E.; Pasqualini, L.; Innocente, S.; Scricciolo, V.; Rignanese, A.; Ciuffetti, G. (1991): Physical training and antiplatelet treatment in stage II peripheral arterial occlusive disease: alone or combined? In: *Angiology* 42 (7), S. 513–521. DOI: 10.1177/000331979104200701.
- Mannion, A. F.; Müntener, M.; Taimela, S.; Dvorak, J. (1999): A randomized clinical trial of three active therapies for chronic low back pain. In: *Spine* 24 (23), S. 2435–2448. DOI: 10.1097/00007632-199912010-00004.
- Marcora (2005): Can progressive resistance training reverse cachexia in patients with rheumatoid arthritis? Results of a pilot study. In: *J Rheumatol* 32, S. 1031.
- MARCUS, BESS H.; SIMKIN, LAUREY R. (1994): The transtheoretical model. In: *Medicine & Science in Sports & Exercise* 26 (11), 1400-1404. DOI: 10.1249/00005768-199411000-00016.
- Marquis, Karine; Debigaré, Richard; Lacasse, Yves; LeBlanc, Pierre; Jobin, Jean; Carrier, Guy; Maltais, François (2002): Mid thigh muscle cross-sectional area is a better predictor of mortality than body mass index in patients with chronic obstructive pulmonary disease. In: *Am J Respir Crit Care Med* 166 (6), S. 809–813. DOI: 10.1164/rccm.2107031.
- Martens, Elisabeth J.; Jonge, Peter de; Na, Beeya; Cohen, Beth E.; Lett, Heather; Whooley, Mary A. (2010): Scared to death? Generalized anxiety disorder and cardiovascular events in patients with stable coronary heart disease: The Heart and Soul Study. In: *Archives of General Psychiatry* 67 (7), S. 750–758. DOI: 10.1001/archgenpsychiatry.2010.74.
- Martinsen (1988): Comparing aerobic and non-aerobic forms of exercise in the treatment of clinical depression: a randomised trial [abstract].
- Martinsen, Egil W.; Hoffart, Asle; Solberg, Øyvind (1989): Comparing aerobic with nonaerobic forms of exercise in the treatment of clinical depression: A randomized trial. In: *Comprehensive Psychiatry* 30 (4), S. 324–331. DOI: 10.1016/0010-440X(89)90057-6.
- Massie, B. M.; Conway, M.; Rajagopalan, B.; Yonge, R.; Frostick, S.; Ledingham, J. et al. (1988): Skeletal muscle metabolism during exercise under ischemic conditions in congestive heart failure. Evidence for abnormalities unrelated to blood flow. In: *Circulation* 78 (2), S. 320–326. DOI: 10.1161/01.CIR.78.2.320.
- Matthews, V. B.; Aström, M-B; Chan, M. H. S.; Bruce, C. R.; Krabbe, K. S.; Prelovsek, O. et al. (2009): Brain-derived neurotrophic factor is produced by skeletal muscle cells in response to contraction and enhances fat oxidation via activation of AMP-activated protein kinase. In: *Diabetologia* 52 (7), S. 1409–1418. DOI: 10.1007/s00125-009-1364-1.
- MCALLISTER, RICHARD M.; HIRAI, TADAKAZU; MUSCH, TIMOTHY I. (1995): Contribution of endothelium-derived nitric oxide (EDNO) to the skeletal muscle blood flow response to exercise. In: *Medicine & Science in Sports & Exercise* 27 (8), 1145-1151. DOI: 10.1249/00005768-199508000-00007.
- McCarthy (2015): Pulmonary rehabilitation for chronic obstructive pulmonary disease. In: *Cochrane Database Syst Rev* 2, CD003793.
- McDonald, C. D.; Burch, G. E.; Walsh, J. J. (1972): Prolonged bed rest in the treatment of idiopathic cardiomyopathy. In: *The American Journal of Medicine* 52 (1), S. 41–50. DOI: 10.1016/0002-9343(72)90006-X.
- McGuire, M. T.; Wing, R. R.; Klem, M. L.; Hill, J. O. (1999): Behavioral strategies of individuals who have maintained long-term weight losses. In: *Obesity Research* 7 (4), S. 334–341. DOI: 10.1002/j.1550-8528.1999.tb00416.x.
- McKay, Heather A.; Petit, Moira A.; Schutz, Robert W.; Prior, Jerilynn C.; Barr, Susan I.; Khan, Karim M. (2000): Augmented trochanteric bone mineral density after modified physical education classes: A randomized school-based exercise intervention study in prepubescent and early pubescent children. In: *The Journal of Pediatrics* 136 (2), S. 156–162. DOI: 10.1016/S0022-3476(00)70095-3.
- McMillan, Elliott M.; Newhouse, Ian J. (2011): Exercise is an effective treatment modality for reducing cancer-related fatigue and improving physical capacity in cancer patients and survivors: a meta-analysis. In: *Applied physiology, nutrition, and metabolism = Physiologie appliquée, nutrition et métabolisme* 36 (6), S. 892–903. DOI: 10.1139/h11-082.
- McNally, P. G.; Watt, P. A.; Rimmer, T.; Burden, A. C.; Hearnshaw, J. R.; Thurston, H. (1994): Impaired contraction and endothelium-dependent relaxation in isolated resistance vessels from patients with insulin-dependent diabetes mellitus. In: *Clinical science (London, England : 1979)* 87 (1), S. 31–36. DOI: 10.1042/cs0870031.
- McNeely, Margaret L.; Campbell, Kristin L.; Rowe, Brian H.; Klassen, Terry P.; Mackey, John R.; Courneya, Kerry S. (2006): Effects of exercise on breast cancer patients and survivors: a systematic review and meta-analysis. In: *Canadian Medical Association Journal* 175 (1), S. 34–41. DOI: 10.1503/cmaj.051073.
- McVeigh (1992): Impaired endothelium-dependent and independent vasodilation in patients with type 2 (non-insulin-dependent) diabetes mellitus. In: *Diabetologia* 35, S. 771.
- Mead (2009): Exercise for depression. In: *Cochrane Database Syst Rev*, CD004366.
- Mehrholz (2010): Treadmill training for patients with Parkinson's disease. In: *Cochrane Database Syst Rev* 20, CD007830.
- Mendes, Felipe A. R.; Gonçalves, Raquel C.; Nunes, Maria P. T.; Saraiva-Romanholo, Beatriz M.; Cukier, Alberto; Stelmach, Rafael et al. (2010): Effects of aerobic training on psychosocial morbidity and symptoms in patients with asthma: a randomized clinical trial. In: *Chest* 138 (2), S. 331–337. DOI: 10.1378/chest.09-2389.

- Metsios, George S.; Stavropoulos-Kalinoglou, Antonios; van Veldhuijzen Zanten, Jet J. C. S.; Nightingale, Peter; Sandoo, Aamer; Dimitroulas, Theodoros et al. (2014): Individualised exercise improves endothelial function in patients with rheumatoid arthritis. In: *Annals of the Rheumatic Diseases* 73 (4), S. 748–751. DOI: 10.1136/annrheumdis-2013-203291.
- Meyer, Katharina; Schwaibold, Matthias; Westbrook, Samuel; Beneke, Ralph; Hajric, Ramiz; Görnandt, Lothar et al. (1996): Effects of short-term exercise training and activity restriction on functional capacity in patients with severe chronic congestive heart failure. In: *The American Journal of Cardiology* 78 (9), S. 1017–1022. DOI: 10.1016/S0002-9149(96)00527-9.
- Meyerhardt, Jeffrey A.; Giovannucci, Edward L.; Holmes, Michelle D.; Chan, Andrew T.; Chan, Jennifer A.; Colditz, Graham A.; Fuchs, Charles S. (2006): Physical activity and survival after colorectal cancer diagnosis. In: *Journal of Clinical Oncology* 24 (22), S. 3527–3534. DOI: 10.1200/JCO.2006.06.0855.
- Meyerhardt, Jeffrey A.; Giovannucci, Edward L.; Ogino, Shuji; Kirkner, Gregory J.; Chan, Andrew T.; Willett, Walter; Fuchs, Charles S. (2009): Physical activity and male colorectal cancer survival. In: *Arch Intern Med* 169 (22), S. 2102–2108. DOI: 10.1001/archinternmed.2009.412.
- Meyerhardt, Jeffrey A.; Heseltine, Denise; Niedzwiecki, Donna; Hollis, Donna; Saltz, Leonard B.; Mayer, Robert J. et al. (2006): Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. In: *Journal of Clinical Oncology* 24 (22), S. 3535–3541. DOI: 10.1200/JCO.2006.06.0863.
- Midtgaard, Julie; STELTER, REINHARD; Rørth, Mikael; Adamsen, Lis (2007): Regaining a sense of agency and shared self-reliance: the experience of advanced disease cancer patients participating in a multidimensional exercise intervention while undergoing chemotherapy--analysis of patient diaries. In: *Scandinavian Journal of Psychology* 48 (2), S. 181–190. DOI: 10.1111/j.1467-9450.2007.00562.x.
- Miedema (1994): Reuma-onderzoek meerdere echelons (ROME): basisrapport.
- Miller, W. C.; Koceja, D. M.; Hamilton, E. J. (1997): A meta-analysis of the past 25 years of weight loss research using diet, exercise or diet plus exercise intervention. In: *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 21 (10), S. 941–947. DOI: 10.1038/sj.ijo.0800499.
- Minor (1988): Exercise tolerance and disease related measures in patients with rheumatoid arthritis and osteoarthritis. In: *J Rheumatol* 15, S. 905.
- Minor, M. A.; Hewett, J. E.; Webel, R. R.; Anderson, S. K.; Kay, D. R. (1989): Efficacy of physical conditioning exercise in patients with rheumatoid arthritis and osteoarthritis. In: *Arthritis and rheumatism* 32 (11), S. 1396–1405. DOI: 10.1002/anr.1780321108.
- Miyai, I.; Fujimoto, Y.; Ueda, Y.; Yamamoto, H.; Nozaki, S.; Saito, T.; Kang, J. (2000): Treadmill training with body weight support: its effect on Parkinson's disease. In: *Archives of Physical Medicine and Rehabilitation* 81 (7), S. 849–852. DOI: 10.1053/apmr.2000.4439.
- Miyai, Ichiro; Fujimoto, Yasuyuki; Yamamoto, Hiroshi; Ueda, Yoshishige; Saito, Toshio; Nozaki, Sonoko; Kang, Jin (2002): Long-term effect of body weight-supported treadmill training in Parkinson's disease: a randomized controlled trial. In: *Archives of Physical Medicine and Rehabilitation* 83 (10), S. 1370–1373. DOI: 10.1053/apmr.2002.34603.
- Moorcroft, A. J.; Dodd, M. E.; Webb, A. K. (1997): Exercise testing and prognosis in adult cystic fibrosis. In: *Thorax* 52 (3), S. 291–293. DOI: 10.1136/thx.52.3.291.
- Motl, R. W.; Gosney, J. L. (2008): Effect of exercise training on quality of life in multiple sclerosis: a meta-analysis. In: *Multiple sclerosis (Houndmills, Basingstoke, England)* 14 (1), S. 129–135. DOI: 10.1177/1352458507080464.
- Mourier, A.; Gautier, J. F.; Kerviler, E. de; Bigard, A. X.; Villette, J. M.; Garnier, J. P. et al. (1997): Mobilization of visceral adipose tissue related to the improvement in insulin sensitivity in response to physical training in NIDDM. Effects of branched-chain amino acid supplements. In: *Diabetes Care* 20 (3), S. 385–391. DOI: 10.2337/diacare.20.3.385.
- Moy, C. S.; Songer, T. J.; LaPorte, R. E.; Dorman, J. S.; Kriska, A. M.; Orchard, T. J. et al. (1993): Insulin-dependent diabetes mellitus, physical activity, and death. In: *American Journal of Epidemiology* 137 (1), S. 74–81. DOI: 10.1093/oxfordjournals.aje.a116604.
- Munkholm, Klaus; PEDERSEN, BENTE KLARLUND; Kessing, Lars Vedel; Vinberg, Maj (2014): Elevated levels of plasma brain derived neurotrophic factor in rapid cycling bipolar disorder patients. In: *Psychoneuroendocrinology* 47, S. 199–211. DOI: 10.1016/j.psychneuen.2014.05.011.
- Myers, Jonathan; Prakash, Manish; Froelicher, Victor; Do, Dat; Partington, Sara; Atwood, J. Edwin (2002): Exercise capacity and mortality among men referred for exercise testing. In: *New England Journal of Medicine* 346 (11), S. 793–801. DOI: 10.1056/NEJMoa011858.
- Mynors-Wallis, L. M.; Gath, D. H.; Day, A.; Baker, F. (2000): Randomised controlled trial of problem solving treatment, antidepressant medication, and combined treatment for major depression in primary care. In: *BMJ* 320 (7226), S. 26–30. DOI: 10.1136/bmj.320.7226.26.
- Naci, Huseyin; Ioannidis, John P. A. (2013): Comparative effectiveness of exercise and drug interventions on mortality outcomes: metaepidemiological study. In: *BMJ* 347 (oct01 1), f5577. DOI: 10.1136/bmj.f5577.
- National Heart, Lung and Blood Institute (1998): Obesity education initiative expert panel: Clinical guidelines on the identification, evaluation and treatment of overweight and obesity in adults: The evidence report.

- National Institutes of Health (1995): Global initiative for asthma.
- National Institutes of Health Consensus Development Panel (1993): Triglyceride, LDL, and CHD. In: *JAMA* 269, S. 505.
- Neuberger, Geri B.; Press, Allan N.; Lindsley, Herbert B.; Hinton, Ruthelystyn; Cagle, Perri E.; Carlson, Katherine et al. (1997): Effects of exercise on fatigue, aerobic fitness, and disease activity measures in persons with rheumatoid arthritis. In: *Res. Nurs. Health* 20 (3), S. 195–204. DOI: 10.1002/(SICI)1098-240X(199706)20:3<195::AID-NUR3>3.0.CO;2-D.
- Nguyen-Duy, Thanh-Binh; Nichaman, Milton Z.; Church, Timothy S.; Blair, Steven N.; Ross, Robert (2003): Visceral fat and liver fat are independent predictors of metabolic risk factors in men. In: *American journal of physiology. Endocrinology and metabolism* 284 (6), E1065–71. DOI: 10.1152/ajpendo.00442.2002.
- Nicklas, Barbara J.; Katzell, Leslie I.; Busby-Whitehead, Jan; Goldberg, Andrew P. (1997): Increases in high-density lipoprotein cholesterol with endurance exercise training are blunted in obese compared with lean men. In: *Metabolism* 46 (5), S. 556–561. DOI: 10.1016/S0026-0495(97)90194-3.
- Nicolaï, Saskia P. A.; Hendriks, Erik J. M.; Prins, Martin H.; Teijink, Joep A. W. (2010): Optimizing supervised exercise therapy for patients with intermittent claudication. In: *Journal of Vascular Surgery* 52 (5), S. 1226–1233. DOI: 10.1016/j.jvs.2010.06.106.
- Nielsen, Paul J.; Hafsdahl, Adam R.; Conn, Vicki S.; LeMaster, Joseph W.; Brown, Sharon A. (2006): Meta-analysis of the effect of exercise interventions on fitness outcomes among adults with type 1 and type 2 diabetes. In: *Diabetes Research and Clinical Practice* 74 (2), S. 111–120. DOI: 10.1016/j.diabres.2006.03.033.
- Nikander, Riku; Sievänen, Harri; Heinonen, Ari; Daly, Robin M.; Uusi-Rasi, Kirsti; Kannus, Pekka (2010): Targeted exercise against osteoporosis: A systematic review and meta-analysis for optimising bone strength throughout life. In: *BMC Medicine* 8 (1), S. 47. DOI: 10.1186/1741-7015-8-47.
- Nixon, P. A.; Orenstein, D. M.; Kelsey, S. F.; Doershuk, C. F. (1992): The prognostic value of exercise testing in patients with cystic fibrosis. In: *The New England journal of medicine* 327 (25), S. 1785–1788. DOI: 10.1056/NEJM199212173272504.
- Nordemar, R.; Edström, L.; Ekblom, B. (1976): Changes in muscle fibre size and physical performance in patients with rheumatoid arthritis after short-term physical training. In: *Scandinavian Journal of Rheumatology* 5 (2), S. 70–76. DOI: 10.3109/03009747609099893.
- Nordesjö, L. O.; Nordgren, B.; Wigren, A.; Kolstad, K. (1983): Isometric strength and endurance in patients with severe rheumatoid arthritis or osteoarthritis in the knee joints. A comparative study in healthy men and women. In: *Scandinavian Journal of Rheumatology* 12 (2), S. 152–156. DOI: 10.3109/03009748309102902.
- Normandin, Edgar A.; McCusker, Corliss; Connors, MaryLou; Vale, Frederick; Gerardi, Daniel; ZuWallack, Richard L. (2002): An evaluation of two approaches to exercise conditioning in pulmonary rehabilitation. In: *Chest* 121 (4), S. 1085–1091. DOI: 10.1378/chest.121.4.1085.
- Norris, Richard; Carroll, Douglas; Cochrane, Raymond (1990): The effects of aerobic and anaerobic training on fitness, blood pressure, and psychological stress and well-being. In: *Journal of Psychosomatic Research* 34 (4), S. 367–375. DOI: 10.1016/0022-3999(90)90060-H.
- Norris, Richard; Carroll, Douglas; Cochrane, Raymond (1992): The effects of physical activity and exercise training on psychological stress and well-being in an adolescent population. In: *Journal of Psychosomatic Research* 36 (1), S. 55–65. DOI: 10.1016/0022-3999(92)90114-H.
- Notarius, C. F.; Azevedo, E. R.; Parker, J. D.; Floras, J. S. (2002): Peak oxygen uptake is not determined by cardiac noradrenaline spillover in heart failure. In: *European Heart Journal* 23 (10), S. 800–805. DOI: 10.1053/euhj.2001.2942.
- Nyberg, Jenny; Åberg, Maria A. I.; Schiöler, Linus; Nilsson, Michael; Wallin, Anders; Torén, Kjell; Kuhn, H. Georg (2014): Cardiovascular and cognitive fitness at age 18 and risk of early-onset dementia. In: *Brain* 137 (Pt 5), S. 1514–1523. DOI: 10.1093/brain/awu041.
- Obrant, K. J.; Bengnér, U.; Johnell, O.; Nilsson, B. E.; Sernbo, I. (1989): Increasing age-adjusted risk of fragility fractures: a sign of increasing osteoporosis in successive generations? In: *Calcified Tissue International* 44 (3), S. 157–167. DOI: 10.1007/BF02556558.
- O'Donovan, G.; Thomas, E. L.; McCarthy, J. P.; Fitzpatrick, J.; Durighel, G.; Mehta, S. et al. (2009): Fat distribution in men of different waist girth, fitness level and exercise habit. In: *International Journal of Obesity* 33 (12), S. 1356–1362. DOI: 10.1038/ijo.2009.189.
- Oesch, Peter; Kool, Jan; Hagen, Kåre Birger; Bachmann, Stefan (2010): Effectiveness of exercise on work disability in patients with non-acute non-specific low back pain: Systematic review and meta-analysis of randomised controlled trials. In: *Journal of Rehabilitation Medicine* 42 (3), S. 193–205. DOI: 10.2340/16501977-0524.
- Ohkawara, K.; Tanaka, S.; Miyachi, M.; Ishikawa-Takata, K.; Tabata, I. (2007): A dose-response relation between aerobic exercise and visceral fat reduction: systematic review of clinical trials. In: *International journal of obesity (2005)* 31 (12), S. 1786–1797. DOI: 10.1038/sj.ijo.0803683.
- Olgiati (1988): Increased energy cost of walking in multiple sclerosis: effect of spasticity, ataxia, and weakness. In: *Arch Phys Med Rehabil* 69, S. 846.

- Olsen, Rasmus H.; Krogh-Madsen, Rikke; Thomsen, Carsten; Booth, Frank W.; Pedersen, Bente K. (2008): Metabolic responses to reduced daily steps in healthy nonexercising men. In: *JAMA* 299 (11), S. 1261–1263. DOI: 10.1001/jama.299.11.1259.
- Olson, Andrea K.; Eadie, Brennan D.; Ernst, Carl; Christie, Brian R. (2006): Environmental enrichment and voluntary exercise massively increase neurogenesis in the adult hippocampus via dissociable pathways. In: *Hippocampus* 16 (3), S. 250–260. DOI: 10.1002/hipo.20157.
- O'Neill, P. A.; Dodds, M.; Phillips, B.; Poole, J.; Webb, A. K. (1987): Regular exercise and reduction of breathlessness in patients with cystic fibrosis. In: *British Journal of Diseases of the Chest* 81, S. 62–69. DOI: 10.1016/0007-0971(87)90109-4.
- Orenstein (1996): The child and the adolescent athlete.
- Orenstein, David M.; Hovell, Melbourne F.; Mulvihill, Mary; Keating, Kristen K.; Hofstetter, C. Richard; Kelsey, Sheryl et al. (2004): Strength vs aerobic training in children with cystic fibrosis: a randomized controlled trial. In: *Chest* 126 (4), S. 1204–1214. DOI: 10.1378/chest.126.4.1204.
- Orozco (2008): Exercise or exercise and diet for preventing type 2 diabetes mellitus. In: *Cochrane Database Syst Rev*, CD003054.
- Orwin, A. (1973): 'The running treatment': a preliminary communication on a new use for an old therapy (physical activity) in the agoraphobic syndrome. In: *The British journal of psychiatry : the journal of mental science* 122 (567), S. 175–179. DOI: 10.1192/bjp.122.2.175.
- Orwin, A. (1974): Treatment of a situational phobia--a case for running. In: *The British journal of psychiatry : the journal of mental science* 125 (0), S. 95–98. DOI: 10.1192/bjp.125.1.95.
- Ott, Susan M.; Aitken, Moira L. (1998): OSTEOPOROSIS IN PATIENTS WITH CYSTIC FIBROSIS. In: *Clinics in Chest Medicine* 19 (3), S. 555–567. DOI: 10.1016/S0272-5231(05)70100-3.
- Owen, A.; Wiles, J.; Swaine, I. (2010): Effect of isometric exercise on resting blood pressure: a meta analysis. In: *Journal of Human Hypertension* 24 (12), S. 796–800. DOI: 10.1038/jhh.2010.13.
- Owens, J. F.; Matthews, K. A.; Wing, R. R.; Kuller, L. H. (1992): Can physical activity mitigate the effects of aging in middle-aged women? In: *Circulation* 85 (4), S. 1265–1270. DOI: 10.1161/01.CIR.85.4.1265.
- Paffenbarger, R. S.; Lee, I. M.; Leung, R. (1994): Physical activity and personal characteristics associated with depression and suicide in American college men. In: *Acta psychiatrica Scandinavica. Supplementum* 377 (s377), S. 16–22. DOI: 10.1111/j.1600-0447.1994.tb05796.x.
- Paganini-Hill, A.; Chao, A.; Ross, R. K.; Henderson, B. E. (1991): Exercise and other factors in the prevention of hip fracture: the Leisure World study. In: *Epidemiology* 2 (1), S. 16–25. DOI: 10.1097/00001648-199101000-00004.
- Pajonk, Frank-Gerald; Wobrock, Thomas; Gruber, Oliver; Scherk, Harald; Berner, Dorothea; Kaizl, Inge et al. (2010): Hippocampal plasticity in response to exercise in schizophrenia. In: *Archives of General Psychiatry* 67 (2), S. 133–143. DOI: 10.1001/archgenpsychiatry.2009.193.
- Palacio, J.; Gáldiz, J. B.; Bech, J. J.; Mariñán, M.; Casadevall, C.; Martínez, P.; Gea, J. (2002): Expresión de los genes de la interleucina 10 (IL-10) y del factor de necrosis tumoral alfa (TNF- $\alpha$ ) en músculos respiratorios y periféricos. Relación con el daño sarcolémico. In: *Archivos de Bronconeumología* 38 (7), S. 311–316. DOI: 10.1016/S0300-2896(02)75224-1.
- Pan, X. R.; Li, G. W.; Hu, Y. H.; Wang, J. X.; Yang, W. Y.; An, Z. X. et al. (1997): Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. In: *Diabetes Care* 20 (4), S. 537–544. DOI: 10.2337/diacare.20.4.537.
- Paolisso, Giuseppe; Riu, Stefano de; Marrazzo, Giuseppe; Verza, Mario; Varricchio, Michele; D'Onofrio, Felice (1991): Insulin resistance and hyperinsulinemia in patients with chronic congestive heart failure. In: *Metabolism* 40 (9), S. 972–977. DOI: 10.1016/0026-0495(91)90075-8.
- Paolucci, Stefano; Gandolfo, Carlo; Provinciali, Leandro; Torta, Riccardo; Toso, Vito (2006): The Italian multicenter observational study on post-stroke depression (DESTRO). In: *Journal of Neurology* 253 (5), S. 556–562. DOI: 10.1007/s00415-006-0058-6.
- PARNELL, Melinda M.; HOLST, Diane P.; KAYE, David M. (2002): Exercise training increases arterial compliance in patients with congestive heart failure. In: *Clinical Science* 102 (1), S. 1–7. DOI: 10.1042/cs1020001.
- Pasco, Julie A.; Williams, Lana J.; Jacka, Felice N.; Henry, Margaret J.; Coulson, Carolyn E.; Brennan, Sharon L. et al. (2011): Habitual physical activity and the risk for depressive and anxiety disorders among older men and women. In: *International Psychogeriatrics* 23 (2), S. 292–298. DOI: 10.1017/S1041610210001833.
- Pasternak (1990): Spectrum of risk factors for CHD. In: *J Am Coll Cardiol* 27, S. 964.
- Pavlou, K. N.; Krey, S.; Steffee, W. P. (1989): Exercise as an adjunct to weight loss and maintenance in moderately obese subjects. In: *The American Journal of Clinical Nutrition* 49 (5 Suppl), S. 1115–1123. DOI: 10.1093/ajcn/49.5.1115.
- Pedersen, Bente K. (2009): The diseaseome of physical inactivity--and the role of myokines in muscle--fat cross talk. In: *The Journal of Physiology* 587 (Pt 23), S. 5559–5568. DOI: 10.1113/jphysiol.2009.179515.
- Pedersen, Bente K. (2011): Exercise-induced myokines and their role in chronic diseases. In: *Brain, Behavior, and Immunity* 25 (5), S. 811–816. DOI: 10.1016/j.bbi.2011.02.010.



- Pedersen, Line; Christensen, Jesper Frank; Hojman, Pernille (2015): Effects of exercise on tumor physiology and metabolism. In: *Cancer journal (Sudbury, Mass.)* 21 (2), S. 111–116. DOI: 10.1097/PPO.0000000000000096.
- Pedersen, Bente K.; FEBBRAIO, MARK A. (2008): Muscle as an endocrine organ: focus on muscle-derived interleukin-6. In: *Physiological Reviews* 88 (4), S. 1379–1406. DOI: 10.1152/physrev.90100.2007.
- Pedersen, Bente K.; Pedersen, Maria; Krabbe, Karen S.; Bruunsgaard, Helle; Matthews, Vance B.; FEBBRAIO, MARK A. (2009): Role of exercise-induced brain-derived neurotrophic factor production in the regulation of energy homeostasis in mammals. In: *Experimental Physiology* 94 (12), S. 1153–1160. DOI: 10.1113/expphysiol.2009.048561.
- Pedersen, B. K.; Saltin, B. (2006): Evidence for prescribing exercise as therapy in chronic disease. In: *Scandinavian journal of medicine & science in sports* 16 Suppl 1 (S1), S. 3–63. DOI: 10.1111/j.1600-0838.2006.00520.x.
- Pedersen, B. K.; Saltin, B. (2015): Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. In: *Scandinavian journal of medicine & science in sports* 25 Suppl 3, S. 1–72. DOI: 10.1111/sms.12581.
- Pedersen, B. K.; Steensberg, A.; Schjerling, P. (2001): Muscle-derived interleukin-6: possible biological effects. In: *The Journal of Physiology* 536 (Pt 2), S. 329–337. DOI: 10.1111/j.1469-7793.2001.0329c.xd.
- Peel, J. Brent; Sui, Xuemei; Matthews, Charles E.; Adams, Swann A.; Hébert, James R.; Hardin, James W. et al. (2009): Cardiorespiratory fitness and digestive cancer mortality: findings from the aerobics center longitudinal study. In: *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 18 (4), S. 1111–1117. DOI: 10.1158/1055-9965.EPI-08-0846.
- Péronnet, F.; Cléroux, J.; Perrault, H.; Cousineau, D.; Champlain, J. de; Nadeau, R. (1981): Plasma norepinephrine response to exercise before and after training in humans. In: *Journal of applied physiology: respiratory, environmental and exercise physiology* 51 (4), S. 812–815. DOI: 10.1152/jappl.1981.51.4.812.
- Perri, Michael G.; McAdoo, William George; McAllister, David A.; Lauer, Joan B.; Yancey, Donna Z. (1986): Enhancing the efficacy of behavior therapy for obesity: Effects of aerobic exercise and a multicomponent maintenance program. In: *Journal of Consulting and Clinical Psychology* 54 (5), S. 670–675. DOI: 10.1037/0022-006X.54.5.670.
- Perri, Michael G.; McAllister, David A.; Gange, James J.; Jordan, Randall C.; McAdoo, W. George; Nezu, Arthur M. (1988): Effects of four maintenance programs on the long-term management of obesity. In: *Journal of Consulting and Clinical Psychology* 56 (4), S. 529–534. DOI: 10.1037/0022-006X.56.4.529.
- Persch, Leslie N.; Ugrinowitsch, Carlos; Pereira, Gleber; Rodacki, André L. F. (2009): Strength training improves fall-related gait kinematics in the elderly: a randomized controlled trial. In: *Clinical biomechanics (Bristol, Avon)* 24 (10), S. 819–825. DOI: 10.1016/j.clinbiomech.2009.07.012.
- Pescatello, Linda S.; Franklin, Barry A.; Fagard, Robert; Farquhar, William B.; Kelley, George A.; Ray, Chester A. (2004): American College of Sports Medicine position stand. Exercise and hypertension. In: *Medicine & Science in Sports & Exercise* 36 (3), S. 533–553. DOI: 10.1249/01.MSS.0000115224.88514.3A.
- Petajan, J. H.; Gappmaier, E.; White, A. T.; Spencer, M. K.; Mino, L.; Hicks, R. W. (1996): Impact of aerobic training on fitness and quality of life in multiple sclerosis. In: *Annals of Neurology* 39 (4), S. 432–441. DOI: 10.1002/ana.410390405.
- Petajan, J. H.; JARCHO, L. W. (1975): Motor unit control in Parkinson's disease and the influence of levodopa. In: *Neurology* 25 (9), S. 866–869. DOI: 10.1212/WNL.25.9.866.
- Petersen, Tom; Kryger, Peter; Ekdahl, C.; Olsen, Steen; Jacobsen, Soren (2002): The effect of McKenzie therapy as compared with that of intensive strengthening training for the treatment of patients with subacute or chronic low back pain: A randomized controlled trial. In: *Spine* 27 (16), S. 1702–1709. DOI: 10.1097/00007632-200208150-00004.
- Petersen, Anne Marie Winther; Magkos, Faidon; Atherton, Philip; Selby, Anna; Smith, Kenneth; Rennie, Michael J. et al. (2007): Smoking impairs muscle protein synthesis and increases the expression of myostatin and MAFbx in muscle. In: *American journal of physiology. Endocrinology and metabolism* 293 (3), E843-8. DOI: 10.1152/ajpendo.00301.2007.
- Petersen, A. M. W.; Mittendorfer, B.; Magkos, F.; Iversen, M.; Pedersen, B. K. (2008): Physical activity counteracts increased whole-body protein breakdown in chronic obstructive pulmonary disease patients. In: *Scandinavian Journal of Medicine and Science in Sports* 18 (5), S. 557–564. DOI: 10.1111/j.1600-0838.2007.00727.x.
- Petersen, Anne Marie W.; PEDERSEN, BENTE KLARLUND (2005): The anti-inflammatory effect of exercise. In: *Journal of applied physiology (Bethesda, Md. : 1985)* 98 (4), S. 1154–1162. DOI: 10.1152/jappphysiol.00164.2004.
- Peurala, Sinikka H.; Karttunen, Auli H.; Sjögren, Tuulikki; Paltamaa, Jaana; Heinonen, Ari (2014): Evidence for the effectiveness of walking training on walking and self-care after stroke: a systematic review and meta-analysis of randomized controlled trials. In: *Journal of Rehabilitation Medicine* 46 (5), S. 387–399. DOI: 10.2340/16501977-1805.
- Piepoli, Massimo F.; Francis, Darrel P. (2004): Exercise training in patients with chronic heart failure: Fig 1. In: *BMJ* 328 (7441), 711.3. DOI: 10.1136/bmj.328.7441.711-b.
- Pinkney, J. H.; Downs, L.; Hopton, M.; Mackness, M. I.; Bolton, C. H. (1999): Endothelial dysfunction in Type 1 diabetes mellitus: relationship with LDL oxidation and the effects of vitamin E. In: *Diabetic Medicine* 16 (12), S. 993–999. DOI: 10.1046/j.1464-5491.1999.00191.x.

- Podewils, Laura Jean; Guallar, Eliseo; Kuller, Lewis H.; Fried, Linda P.; Lopez, Oscar L.; Carlson, Michelle; Lyketsos, Constantine G. (2005): Physical activity, APOE genotype, and dementia risk: findings from the Cardiovascular Health Cognition Study. In: *American Journal of Epidemiology* 161 (7), S. 639–651. DOI: 10.1093/aje/kwi092.
- Polese, Janaine C.; Ada, Louise; Dean, Catherine M.; Nascimento, Lucas R.; Teixeira-Salmela, Luci F. (2013): Treadmill training is effective for ambulatory adults with stroke: a systematic review. In: *Journal of Physiotherapy* 59 (2), S. 73–80. DOI: 10.1016/S1836-9553(13)70159-0.
- Pradhan, A. D.; Manson, J. E.; Rifai, N.; Buring, J. E.; Ridker, P. M. (2001): C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. In: *JAMA* 286 (3), S. 327–334. DOI: 10.1001/jama.286.3.327.
- Pronk, N. P. (1993): Short term effects of exercise on plasma lipids and lipoproteins in humans. In: *Sports Medicine* 16 (6), S. 431–448. DOI: 10.2165/00007256-199316060-00006.
- Province, Michael A. (1995): The Effects of Exercise on Falls in Elderly Patients. In: *JAMA* 273 (17), S. 1341. DOI: 10.1001/jama.1995.03520410035023.
- Quirk, H.; Blake, H.; Tennyson, R.; Randell, T. L.; Glazebrook, C. (2014): Physical activity interventions in children and young people with Type 1 diabetes mellitus: a systematic review with meta-analysis. In: *Diabetic medicine : a journal of the British Diabetic Association* 31 (10), S. 1163–1173. DOI: 10.1111/dme.12531.
- Rabasa-Lhoret, R.; Bourque, J.; Ducros, F.; Chiasson, J. L. (2001): Guidelines for premeal insulin dose reduction for postprandial exercise of different intensities and durations in type 1 diabetic subjects treated intensively with a basal-bolus insulin regimen (ultralente-lispro). In: *Diabetes Care* 24 (4), S. 625–630. DOI: 10.2337/diacare.24.4.625.
- Raglin (1997): Physical activity and mental health.
- Ragnarsson (1988): Physiologic effects of functional electrical stimulation-induced exercises in spinal cord-injured individuals. In: *Clin Orthop* 000, S. 53.
- Ram, F. S.; Robinson, S. M.; Black, P. N. (2000): Effects of physical training in asthma: a systematic review. In: *British Journal of Sports Medicine* 34 (3), S. 162–167. DOI: 10.1136/bjsm.34.3.162.
- Rand, Debbie; Eng, Janice J.; Tang, Pei-Fang; Jeng, Jiann-Shing; Hung, Chihya (2009): How active are people with stroke?: use of accelerometers to assess physical activity. In: *Stroke* 40 (1), S. 163–168. DOI: 10.1161/STROKEAHA.108.523621.
- Rastad, Cecilia; Martin, Cathrin; Asenlöf, Pernilla (2014): Barriers, benefits, and strategies for physical activity in patients with schizophrenia. In: *Physical Therapy* 94 (10), S. 1467–1479. DOI: 10.2522/ptj.20120443.
- Ravaglia, G.; Forti, P.; Lucicesare, A.; Pisacane, N.; Rietti, E.; Bianchin, M.; Dalmonte, E. (2008): Physical activity and dementia risk in the elderly: findings from a prospective Italian study. In: *Neurology* 70 (19 Pt 2), S. 1786–1794. DOI: 10.1212/01.wnl.0000296276.50595.86.
- Recommendations for exercise training in chronic heart failure patients (2001). In: *European Heart Journal* 22 (2), S. 125–135.
- Regensteiner, Judith G.; Steiner, John F.; Hiatt, William R. (1996): Exercise training improves functional status in patients with peripheral arterial disease. In: *Journal of Vascular Surgery* 23 (1), S. 104–115. DOI: 10.1016/S0741-5214(05)80040-0.
- Regensteiner, Judith G.; Ware, John E.; McCarthy, Walter J.; Zhang, Peter; Forbes, William P.; Heckman, Jeffrey; Hiatt, William R. (2002): Effect of cilostazol on treadmill walking, community-based walking ability, and health-related quality of life in patients with intermittent claudication due to peripheral arterial disease: meta-analysis of six randomized controlled trials. In: *Journal of the American Geriatrics Society* 50 (12), S. 1939–1946. DOI: 10.1046/j.1532-5415.2002.50604.x.
- Reitman, J. S.; Vasquez, B.; Klimes, I.; Nagulesparan, M. (1984): Improvement of glucose homeostasis after exercise training in non-insulin-dependent diabetes. In: *Diabetes Care* 7 (5), S. 434–441. DOI: 10.2337/diacare.7.5.434.
- Rietberg (2005): Exercise therapy for multiple sclerosis. In: *Cochrane Database Syst Rev*, CD003980.
- Riggs, A. C.; Seaquist, E. R.; Moran, A. (1999): Guidelines for the diagnosis and therapy of diabetes mellitus in cystic fibrosis. In: *Current Opinion in Pulmonary Medicine* 5 (6), S. 378–382. DOI: 10.1097/00063198-199911000-00010.
- Rimmele, Ulrike; Zellweger, Bea Costa; Marti, Bernard; Seiler, Roland; Mohiyeddini, Changiz; Ehlert, Ulrike; Heinrichs, Markus (2007): Trained men show lower cortisol, heart rate and psychological responses to psychosocial stress compared with untrained men. In: *Psychoneuroendocrinology* 32 (6), S. 627–635. DOI: 10.1016/j.psyneuen.2007.04.005.
- Rissanen (1991): Determinants of weight gain and overweight in adult Finns. In: *Eur J Clin Nutr* 45, S. 419.
- Robertson, M. Clare; Campbell, A. John; Gardner, Melinda M.; Devlin, Nancy (2002): Preventing injuries in older people by preventing falls: a meta-analysis of individual-level data. In: *Journal of the American Geriatrics Society* 50 (5), S. 905–911. DOI: 10.1046/j.1532-5415.2002.50218.x.
- Robillon (1994): Abnormalities suggestive of cardiomyopathy in patients with type 2 diabetes of relatively short duration. In: *Diabetes Metab* 20, S. 473.
- Rodgers (1999): Gait characteristics of individuals with multiple sclerosis before and after a 6-month aerobic training program. In: *J Rehabil Res Dev* 36, S. 183.

- Rogers, M. W.; Probst, M. M.; Gruber, J. J.; Berger, R.; Boone, J. B. (1996): Differential effects of exercise training intensity on blood pressure and cardiovascular responses to stress in borderline hypertensive humans. In: *Journal of Hypertension* 14 (11), S. 1369–1375. DOI: 10.1097/00004872-199611000-00017.
- Rolland, Yves; Pillard, Fabien; Klapouszczak, Adrian; Reynish, Emma; Thomas, David; Andrieu, Sandrine et al. (2007): Exercise program for nursing home residents with Alzheimer's disease: a 1-year randomized, controlled trial. In: *Journal of the American Geriatrics Society* 55 (2), S. 158–165. DOI: 10.1111/j.1532-5415.2007.01035.x.
- Rönnemaa, T.; Mattila, K.; Lehtonen, A.; Kallio, V. (1986): A controlled randomized study on the effect of long-term physical exercise on the metabolic control in type 2 diabetic patients. In: *Acta Medica Scandinavica* 220 (3), S. 219–224. DOI: 10.1111/j.0954-6820.1986.tb02754.x.
- Roos, Ewa M.; Dahlberg, Leif (2005): Positive effects of moderate exercise on glycosaminoglycan content in knee cartilage: a four-month, randomized, controlled trial in patients at risk of osteoarthritis. In: *Arthritis and rheumatism* 52 (11), S. 3507–3514. DOI: 10.1002/art.21415.
- Roos, Ewa M.; Herzog, Walter; Block, Joel A.; Bennell, Kim L. (2011): Muscle weakness, afferent sensory dysfunction and exercise in knee osteoarthritis. In: *Nature Reviews Rheumatology* 7 (1), S. 57–63. DOI: 10.1038/nrrheum.2010.195.
- Rosendahl, Erik; Gustafson, Yngve; Nordin, Ellinor; Lundin-Olsson, Lillemor; Nyberg, Lars (2008): A randomized controlled trial of fall prevention by a high-intensity functional exercise program for older people living in residential care facilities. In: *Aging Clinical and Experimental Research* 20 (1), S. 67–75. DOI: 10.1007/BF03324750.
- Ross, Robert; Aru, James; Freeman, Jennifer; Hudson, Robert; Janssen, Ian (2002): Abdominal adiposity and insulin resistance in obese men. In: *American journal of physiology. Endocrinology and metabolism* 282 (3), E657–63. DOI: 10.1152/ajpendo.00469.2001.
- Ross, R.; Dagnone, D.; Jones, P. J.; Smith, H.; Paddags, A.; Hudson, R.; Janssen, I. (2000): Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. In: *Annals of Internal Medicine* 133 (2), S. 92–103. DOI: 10.7326/0003-4819-133-2-200007180-00008.
- Ross, R.; Janssen, I. (2001): Physical activity, total and regional obesity: dose-response considerations. In: *Medicine & Science in Sports & Exercise* 33 (6 Suppl), S521–7; discussion S528–9. DOI: 10.1097/00005768-200106001-00023.
- Ross, Robert; Janssen, Ian; Dawson, Jody; Kungl, Ann-Marie; Kuk, Jennifer L.; Wong, Suzy L. et al. (2004): Exercise-induced reduction in obesity and insulin resistance in women: a randomized controlled trial. In: *Obesity Research* 12 (5), S. 789–798. DOI: 10.1038/oby.2004.95.
- Rovio, Suvi; Kåreholt, Ingemar; Helkala, Eeva-Liisa; Viitanen, Matti; Winblad, Bengt; Tuomilehto, Jaakko et al. (2005): Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. In: *The Lancet Neurology* 4 (11), S. 705–711. DOI: 10.1016/S1474-4422(05)70198-8.
- Rovio, Suvi; Kåreholt, Ingemar; Viitanen, Matti; Winblad, Bengt; Tuomilehto, Jaakko; Soininen, Hilka et al. (2007): Work-related physical activity and the risk of dementia and Alzheimer's disease. In: *International Journal of Geriatric Psychiatry* 22 (9), S. 874–882. DOI: 10.1002/gps.1755.
- Ruderman, N. B.; Ganda, O. P.; Johansen, K. (1979): The effect of physical training on glucose tolerance and plasma lipids in maturity-onset diabetes. In: *Diabetes* 28 Suppl 1 (Supplement\_1), S. 89–92. DOI: 10.2337/diab.28.1.S89.
- Rugbjerg, Mette; Iepsen, Ulrik Winning; Jørgensen, Karsten Juhl; Lange, Peter (2015): Effectiveness of pulmonary rehabilitation in COPD with mild symptoms: a systematic review with meta-analyses. In: *International Journal of Chronic Obstructive Pulmonary Disease* 10, S. 791–801. DOI: 10.2147/COPD.S78607.
- Ryan, Alice S.; Ivey, Frederick M.; Prior, Steven; Li, Guoyan; Hafer-Macko, Charlene (2011): Skeletal muscle hypertrophy and muscle myostatin reduction after resistive training in stroke survivors. In: *Stroke* 42 (2), S. 416–420. DOI: 10.1161/STROKEAHA.110.602441.
- Salman, Ghassan F.; Mosier, Michael C.; Beasley, Brent W.; Calkins, David R. (2003): Rehabilitation for patients with chronic obstructive pulmonary disease: meta-analysis of randomized controlled trials. In: *Journal of General Internal Medicine* 18 (3), S. 213–221. DOI: 10.1046/j.1525-1497.2003.20221.x.
- Salminen (2009): Effects of risk-based multifactorial fall prevention on postural balance in the community-dwelling aged: a randomized controlled trial. In: *Arch Gerontol Geriatr* 48, S. 22.
- Salmon, Peter (2001): Effects of physical exercise on anxiety, depression, and sensitivity to stress. In: *Clinical Psychology Review* 21 (1), S. 33–61. DOI: 10.1016/S0272-7358(99)00032-X.
- Saltin (2000): [Metabolic capacity of skeletal muscles and health]. In: *Ugeskr Laeger* 162, S. 2159.
- Saltin, B.; Henriksson, J.; Nygaard, E.; Andersen, P.; Jansson, E. (1977): Fiber types and metabolic potentials of skeletal muscles in sedentary man and endurance runners. In: *Annals of the New York Academy of Sciences* 301 (1 The Marathon), S. 3–29. DOI: 10.1111/j.1749-6632.1977.tb38182.x.

- Samad, A. K. A.; Taylor, R. S.; Marshall, T.; Chapman, M. A. S. (2005): A meta-analysis of the association of physical activity with reduced risk of colorectal cancer. In: *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland* 7 (3), S. 204–213. DOI: 10.1111/j.1463-1318.2005.00747.x.
- Sandanger (2000): Relation between health problems and sickness absence: gender and age differences - a comparison of low-back pain, psychiatric disorders, and injuries. In: *Scand J Public Health* 28, S. 244.
- Sanford-Smith (1998): Therapeutic benefit of aquaerobics for individuals with rheumatoid arthritis. In: *Physiotherapy Canada* Winter, S. 40.
- Santuz, P.; Baraldi, E.; Filippone, M.; Zacchello, F. (1997): Exercise performance in children with asthma: is it different from that of healthy controls? In: *European Respiratory Journal* 10 (6), S. 1254–1260. DOI: 10.1183/09031936.97.10061254.
- Sarlio-Lähteenkorva, S.; Rissanen, A. (1998): Weight loss maintenance: determinants of long-term success. In: *Eating and weight disorders : EWD* 3 (3), S. 131–135. DOI: 10.1007/BF03340000.
- Sarlio-Lähteenkorva, S.; Rissanen, A.; Kaprio, J. (2000): A descriptive study of weight loss maintenance: 6 and 15 year follow-up of initially overweight adults. In: *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 24 (1), S. 116–125. DOI: 10.1038/sj.ijo.0801094.
- Saunders (2013): Physical fitness training for stroke patients. In: *Cochrane Database Syst Rev* 10, CD003316.
- Scarmeas, Nikolaos; Luchsinger, Jose A.; Schupf, Nicole; Brickman, Adam M.; Cosentino, Stephanie; Tang, Ming X.; Stern, Yaakov (2009): Physical activity, diet, and risk of Alzheimer disease. In: *JAMA* 302 (6), S. 627–637. DOI: 10.1001/jama.2009.1144.
- Schaafsma (2010): Physical conditioning programs for improving work outcomes in workers with back pain. In: *Cochrane Database Syst Rev* 20, CD001822.
- Scheewe, Thomas W.; van Haren, Neeltje E. M.; Sarkisyan, Gayane; Schnack, Hugo G.; Brouwer, Rachel M.; Glint, Maria de et al. (2013): Exercise therapy, cardiorespiratory fitness and their effect on brain volumes: a randomised controlled trial in patients with schizophrenia and healthy controls. In: *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology* 23 (7), S. 675–685. DOI: 10.1016/j.euroneuro.2012.08.008.
- Schenkman, Margaret; Hall, Deborah A.; Barón, Anna E.; Schwartz, Robert S.; Mettler, Pamela; Kohrt, Wendy M. (2012): Exercise for people in early- or mid-stage Parkinson disease: a 16-month randomized controlled trial. In: *Physical Therapy* 92 (11), S. 1395–1410. DOI: 10.2522/ptj.20110472.
- Scherer, T. A.; Spengler, C. M.; Owassapian, D.; Imhof, E.; Boutellier, U. (2000): Respiratory muscle endurance training in chronic obstructive pulmonary disease: impact on exercise capacity, dyspnea, and quality of life. In: *Am J Respir Crit Care Med* 162 (5), S. 1709–1714. DOI: 10.1164/ajrccm.162.5.9912026.
- Schmitz, Kathryn H.; Courneya, Kerry S.; Matthews, Charles; Demark-Wahnefried, Wendy; GALVÃO, DANIEL A.; Pinto, Bernardine M. et al. (2010): American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. In: *Medicine and science in sports and exercise* 42 (7), S. 1409–1426. DOI: 10.1249/MSS.0b013e3181e0c112.
- Schneider, S. H.; Amorosa, L. F.; Khachaturian, A. K.; Ruderman, N. B. (1984): Studies on the mechanism of improved glucose control during regular exercise in type 2 (non-insulin-dependent) diabetes. In: *Diabetologia* 26 (5), S. 355–360. DOI: 10.1007/BF00266036.
- Schneiderman-Walker, J.; Pollock, S. L.; Corey, M.; Wilkes, D. D.; Canny, G. J.; Pedder, L.; Reisman, J. J. (2000): A randomized controlled trial of a 3-year home exercise program in cystic fibrosis. In: *The Journal of Pediatrics* 136 (3), S. 304–310. DOI: 10.1067/mpd.2000.103408.
- Schoepf, D.; Potluri, R.; Uppal, H.; Natalwala, A.; Narendran, P.; Heun, R. (2012): Type-2 diabetes mellitus in schizophrenia: increased prevalence and major risk factor of excess mortality in a naturalistic 7-year follow-up. In: *European Psychiatry* 27 (1), S. 33–42. DOI: 10.1016/j.eurpsy.2011.02.009.
- Schonstein (2003): Work conditioning, work hardening and functional restoration for workers with back and neck pain. In: *Cochrane Database Syst Rev*, CD001822.
- Schuit, A. J.; Feskens, E. J.; Launer, L. J.; Kromhout, D. (2001): Physical activity and cognitive decline, the role of the apolipoprotein e4 allele. In: *Medicine & Science in Sports & Exercise* 33 (5), S. 772–777. DOI: 10.1097/00005768-200105000-00015.
- Schulze, Paul Christian; Gielen, Stephan; Schuler, Gerhard; Hambrecht, Rainer (2002): Chronic heart failure and skeletal muscle catabolism: effects of exercise training. In: *International Journal of Cardiology* 85 (1), S. 141–149. DOI: 10.1016/S0167-5273(02)00243-7.
- Scott (1960): The contributions of physical activity to psychological development. In: *Res Q* 31, S. 307.
- Seeger, J. P. H.; Thijssen, D. H. J.; Noordam, K.; Cranen, M. E. C.; Hopman, M. T. E.; Nijhuis-van der Sanden, M. W. G. (2011): Exercise training improves physical fitness and vascular function in children with type 1 diabetes. In: *Diabetes, obesity & metabolism* 13 (4), S. 382–384. DOI: 10.1111/j.1463-1326.2011.01361.x.
- Segal, Neil A.; Glass, Natalie A. (2011): Is quadriceps muscle weakness a risk factor for incident or progressive knee osteoarthritis? In: *The Physician and Sportsmedicine* 39 (4), S. 44–50. DOI: 10.3810/psm.2011.11.1938.

- Sell (1994): The effect of physical training on psychiatric patients.
- Selvadurai, H. C.; Blimkie, C. J.; Meyers, N.; Mellis, C. M.; Cooper, P. J.; van Asperen, P. P. (2002): Randomized controlled study of in-hospital exercise training programs in children with cystic fibrosis. In: *Pediatric Pulmonology* 33 (3), S. 194–200. DOI: 10.1002/ppul.10015.
- Semanik, Pamela A.; Chang, Rowland W.; Dunlop, Dorothy D. (2012): Aerobic activity in prevention and symptom control of osteoarthritis. In: *PM & R : the journal of injury, function, and rehabilitation* 4 (5 Suppl), S37–44. DOI: 10.1016/j.pmrj.2012.02.015.
- Sexton, H.; Maere, A.; Dahl, N. H. (1989): Exercise intensity and reduction in neurotic symptoms. A controlled follow-up study. In: *Acta Psychiatrica Scandinavica* 80 (3), S. 231–235. DOI: 10.1111/j.1600-0447.1989.tb01332.x.
- Shaban (2014): The effects of a 2 week modified high intensity interval training program on the homeostatic model of insulin resistance (HOMA-IR) in adults with type 2 diabetes. In: *J Sports Med Phys Fitness* 54, S. 203.
- Shalhoub, Joseph; Qureshi, Mahim; Davies, Alun (2009): Supervised exercise in intermittent claudication: a sedentary notion? In: *Vascular* 17 (2), S. 66–73. DOI: 10.2310/6670.2009.00005.
- Sharif, Salaheddin; Thomas, James M.; Donley, David A.; Gilleland, Diana L.; Bonner, Daniel E.; McCrory, Jean L. et al. (2011): Resistance exercise reduces skeletal muscle cachexia and improves muscle function in rheumatoid arthritis. In: *Case Reports in Medicine* 2011, S. 205691. DOI: 10.1155/2011/205691.
- Shaw (2006): Exercise for overweight or obesity. In: *Cochrane Database Syst Rev*, CD003817.
- Shergill, Sukhwinder S.; Murray, Robin M.; McGuire, Philip K. (1998): Auditory hallucinations: a review of psychological treatments. In: *Schizophrenia Research* 32 (3), S. 137–150. DOI: 10.1016/S0920-9964(98)00052-8.
- Shigematsu, Ryosuke; Okura, Tomohiro; Nakagaichi, Masaki; Tanaka, Kiyoji; Sakai, Tomoaki; Kitazumi, Suguru; Rantanen, Taina (2008): Square-stepping exercise and fall risk factors in older adults: a single-blind, randomized controlled trial. In: *The Journals of Gerontology: Series A* 63 (1), S. 76–82. DOI: 10.1093/gerona/63.1.76.
- Shigematsu, Ryosuke; Okura, Tomohiro; Sakai, Tomoaki; Rantanen, Taina (2008): Square-stepping exercise versus strength and balance training for fall risk factors. In: *Aging Clinical and Experimental Research* 20 (1), S. 19–24. DOI: 10.1007/BF03324743.
- Shirazi, Kambiz Karimzadeh; Wallace, Louise M.; Niknami, Shamsaddin; Hidarnia, Alireza; Torkaman, Giti; Gilchrist, Mollie; Faghihzadeh, Soghrate (2007): A home-based, transtheoretical change model designed strength training intervention to increase exercise to prevent osteoporosis in Iranian women aged 40–65 years: a randomized controlled trial. In: *Health Education Research* 22 (3), S. 305–317. DOI: 10.1093/her/cyl067.
- Shojaee-Moradie, F.; Baynes, K. C. R.; Pentecost, C.; Bell, J. D.; Thomas, E. L.; Jackson, N. C. et al. (2007): Exercise training reduces fatty acid availability and improves the insulin sensitivity of glucose metabolism. In: *Diabetologia* 50 (2), S. 404–413. DOI: 10.1007/s00125-006-0498-7.
- Sigal, Ronald J.; Kenny, Glen P.; Wasserman, David H.; Castaneda-Sceppa, Carmen (2004): Physical activity/exercise and type 2 diabetes. In: *Diabetes Care* 27 (10), S. 2518–2539. DOI: 10.2337/diacare.27.10.2518.
- Sikand (1988): Two-year follow-up of patients treated with a very-low-calorie diet and exercise training. In: *J Am Diet Assoc* 88, S. 487.
- Silva, R. A.; Vieira, R. P.; Duarte, A. C. S.; Lopes, F. D. T. Q. S.; Perini, A.; Mauad, T. et al. (2010): Aerobic training reverses airway inflammation and remodelling in an asthma murine model. In: *The European respiratory journal* 35 (5), S. 994–1002. DOI: 10.1183/09031936.00049509.
- Skelton, D. A.; Beyer, N. (2003): Exercise and injury prevention in older people. In: *Scandinavian journal of medicine & science in sports* 13 (1), S. 77–85. DOI: 10.1034/j.1600-0838.2003.00300.x.
- Skyrme-Jones, R. Andrew P.; O'Brien, Richard C.; Luo, Ming; Meredith, Ian T. (2000): Endothelial vasodilator function is related to low-density lipoprotein particle size and low-density lipoprotein vitamin E content in type 1 diabetes. In: *Journal of the American College of Cardiology* 35 (2), S. 292–299. DOI: 10.1016/S0735-1097(99)00547-1.
- Slemenda, Charles; Heilman, Douglas K.; Brandt, Kenneth D.; Katz, Barry P.; Mazucca, Steven A.; Braunstein, Ethan M.; Byrd, Donna (1998): Reduced quadriceps strength relative to body weight: A risk factor for knee osteoarthritis in women? In: *Arthritis & Rheumatism* 41 (11), S. 1951–1959. DOI: 10.1002/1529-0131(199811)41:11<1951::AID-ART9>3.0.CO;2-9.
- Smeltzer, Suzanne C.; Levietes, Marc H.; Cook, Stuart D. (1996): Expiratory training in multiple sclerosis. In: *Archives of Physical Medicine and Rehabilitation* 77 (9), S. 909–912. DOI: 10.1016/S0003-9993(96)90281-6.
- Smith, Robyn M.; Adeney-Steel, Marion; Fulcher, Gary; Longley, Wendy A. (2006): Symptom change with exercise is a temporary phenomenon for people with multiple sclerosis. In: *Archives of Physical Medicine and Rehabilitation* 87 (5), S. 723–727. DOI: 10.1016/j.apmr.2006.01.015.
- Smith, Ashley Wilder; Alfano, Catherine M.; Reeve, Bryce B.; Irwin, Melinda L.; Bernstein, Leslie; Baumgartner, Kathy et al. (2009): Race/ethnicity, physical activity, and quality of life in breast cancer survivors. In: *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 18 (2), S. 656–663. DOI: 10.1158/1055-9965.EPI-08-0352.

- Smith, S. C.; Blair, S. N.; Bonow, R. O.; Brass, L. M.; Cerqueira, M. D.; Dracup, K. et al. (2001): AHA/ACC Scientific Statement: AHA/ACC guidelines for preventing heart attack and death in patients with atherosclerotic cardiovascular disease: 2001 update: A statement for healthcare professionals from the American Heart Association and the American College of Cardiology. In: *Circulation* 104 (13), S. 1577–1579. DOI: 10.1161/hc3801.097475.
- Smits, P.; Kapma, J. A.; Jacobs, M. C.; Lutterman, J.; Thien, T. (1993): Endothelium-dependent vascular relaxation in patients with type I diabetes. In: *Diabetes* 42 (1), S. 148–153. DOI: 10.2337/diab.42.1.148.
- Snook, Erin M.; Motl, Robert W. (2009): Effect of exercise training on walking mobility in multiple sclerosis: a meta-analysis. In: *Neurorehabilitation and Neural Repair* 23 (2), S. 108–116. DOI: 10.1177/1545968308320641.
- Snowling, Neil J.; Hopkins, Will G. (2006): Effects of different modes of exercise training on glucose control and risk factors for complications in type 2 diabetic patients: a meta-analysis. In: *Diabetes Care* 29 (11), S. 2518–2527. DOI: 10.2337/dc06-1317.
- Socialstyrelsen (2000): Nationella riktlinjer för strokesjukvård.
- Solal, Alain Cohen; Chabernaud, Jean Marc; Gourgon, René (1990): Comparison of oxygen uptake during bicycle exercise in patients with chronic heart failure and in normal subjects. In: *Journal of the American College of Cardiology* 16 (1), S. 80–85. DOI: 10.1016/0735-1097(90)90460-7.
- Soo, K.; Furler, S. M.; Samaras, K.; Jenkins, A. B.; Campbell, L. V.; Chisholm, D. J. (1996): Glycemic responses to exercise in IDDM after simple and complex carbohydrate supplementation. In: *Diabetes Care* 19 (6), S. 575–579. DOI: 10.2337/diacare.19.6.575.
- Speelman, Arlène D.; van de Warrenburg, Bart P.; van Nimwegen, Marlies; Petzinger, Giselle M.; Munneke, Marten; Bloem, Bastiaan R. (2011): How might physical activity benefit patients with Parkinson disease? In: *Nature Reviews Neurology* 7 (9), S. 528–534. DOI: 10.1038/nrneuro.2011.107.
- Sprung, Victoria S.; Cuthbertson, Daniel J.; Pugh, Christopher J. A.; Aziz, Nabil; Kemp, Graham J.; Daousi, Christina et al. (2013): Exercise training in polycystic ovarian syndrome enhances flow-mediated dilation in the absence of changes in fatness. In: *Medicine and science in sports and exercise* 45 (12), S. 2234–2242. DOI: 10.1249/MSS.0b013e31829ba9a1.
- Srikanthan, Preethi; Korenman, Stanley; Davis, Susan (2006): Polycystic ovarian syndrome: the next cardiovascular dilemma in women? In: *Endocrinology and Metabolism Clinics of North America* 35 (3), 611–31, x. DOI: 10.1016/j.ecl.2006.05.001.
- Stamler, J.; Vaccaro, O.; Neaton, J. D.; Wentworth, D. (1993): Diabetes, other risk factors, and 12-yr cardiovascular mortality for men screened in the Multiple Risk Factor Intervention Trial. In: *Diabetes Care* 16 (2), S. 434–444. DOI: 10.2337/diacare.16.2.434.
- Stavropoulos-Kalinoglou, Antonios; Metsios, Giorgos S.; van Veldhuijzen Zanten, Jet J. J. C. S.; Nightingale, Peter; Kitis, George D.; Koutedakis, Yiannis (2013): Individualised aerobic and resistance exercise training improves cardiorespiratory fitness and reduces cardiovascular risk in patients with rheumatoid arthritis. In: *Annals of the Rheumatic Diseases* 72 (11), S. 1819–1825. DOI: 10.1136/annrheumdis-2012-202075.
- Steen, R. Grant; Mull, Courtney; McClure, Robert; Hamer, Robert M.; Lieberman, Jeffrey A. (2006): Brain volume in first-episode schizophrenia: systematic review and meta-analysis of magnetic resonance imaging studies. In: *The British journal of psychiatry: the journal of mental science* 188 (6), S. 510–518. DOI: 10.1192/bjp.188.6.510.
- Stefanick (1994): Physical activity, fitness, health. International. Proceedings and consensus statement.
- Stefanick, M. L.; Mackey, S.; Sheehan, M.; Ellsworth, N.; Haskell, W. L.; Wood, P. D. (1998): Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol. In: *The New England journal of medicine* 339 (1), S. 12–20. DOI: 10.1056/NEJM199807023390103.
- Steinberg, Martin; Leoutsakos, Jeannie-Marie Sheppard; Podewils, Laura Jean; Lyketsos, C. G. (2009): Evaluation of a home-based exercise program in the treatment of Alzheimer's disease: the Maximizing Independence in Dementia (MIND) study. In: *International Journal of Geriatric Psychiatry* 24 (7), S. 680–685. DOI: 10.1002/gps.2175.
- Stewart (2001): ACSM's resource manual for guidelines for exercise testing and prescription.
- Stewart, Kerry J. (2002): Exercise training and the cardiovascular consequences of type 2 diabetes and hypertension: plausible mechanisms for improving cardiovascular health. In: *JAMA* 288 (13), S. 1622–1631. DOI: 10.1001/jama.288.13.1622.
- Strasser, Barbara; Leeb, Gunther; Strehlow, Christoph; Schobersberger, Wolfgang; Haber, Paul; Cauza, Edmund (2011): The effects of strength and endurance training in patients with rheumatoid arthritis. In: *Clinical Rheumatology* 30 (5), S. 623–632. DOI: 10.1007/s10067-010-1584-2.
- Strasser, Barbara; Siebert, Uwe; Schobersberger, Wolfgang (2010): Resistance training in the treatment of the metabolic syndrome: a systematic review and meta-analysis of the effect of resistance training on metabolic clustering in patients with abnormal glucose metabolism. In: *Sports medicine (Auckland, N.Z.)* 40 (5), S. 397–415. DOI: 10.2165/11531380-000000000-00000.
- Sui, Xuemei; Laditka, James N.; Church, Timothy S.; Hardin, James W.; Chase, Nancy; Davis, Keith; Blair, Steven N. (2009): Prospective study of cardiorespiratory fitness and depressive symptoms in women and men. In: *Journal of Psychiatric Research* 43 (5), S. 546–552. DOI: 10.1016/j.jpsychires.2008.08.002.
- Sui, Xuemei; LaMonte, Michael J.; Blair, Steven N. (2007): Cardiorespiratory fitness as a predictor of nonfatal cardiovascular events in asymptomatic women and men. In: *American Journal of Epidemiology* 165 (12), S. 1413–1423. DOI: 10.1093/aje/kwm031.

- Sullivan, M. J.; Green, H. J.; Cobb, F. R. (1990): Skeletal muscle biochemistry and histology in ambulatory patients with long-term heart failure. In: *Circulation* 81 (2), S. 518–527. DOI: 10.1161/01.CIR.81.2.518.
- Sullivan, M. J.; Higginbotham, M. B.; Cobb, F. R. (1988): Exercise training in patients with severe left ventricular dysfunction. Hemodynamic and metabolic effects. In: *Circulation* 78 (3), S. 506–515. DOI: 10.1161/01.CIR.78.3.506.
- Sullivan, M. J.; Higginbotham, M. B.; Cobb, F. R. (1989): Exercise training in patients with chronic heart failure delays ventilatory anaerobic threshold and improves submaximal exercise performance. In: *Circulation* 79 (2), S. 324–329. DOI: 10.1161/01.CIR.79.2.324.
- Sullivan, M. J.; Knight, J. D.; Higginbotham, M. B.; Cobb, F. R. (1989): Relation between central and peripheral hemodynamics during exercise in patients with chronic heart failure. Muscle blood flow is reduced with maintenance of arterial perfusion pressure. In: *Circulation* 80 (4), S. 769–781. DOI: 10.1161/01.CIR.80.4.769.
- Svendsen (1994): Six months' follow-up on exercise added to a short-term diet in overweight postmenopausal women - effects on body composition, resting metabolic rate, cardiovascular risk factors and bone. In: *Int J Obes Relat Metab Disord* 18, S. 692.
- Swedberg, Karl; Cleland, John; Dargie, Henry; Drexler, Helmut; Follath, Ferenc; Komajda, Michel et al. (2005): Guidelines for the diagnosis and treatment of chronic heart failure: executive summary (update 2005): The Task Force for the Diagnosis and Treatment of Chronic Heart Failure of the European Society of Cardiology. In: *European Heart Journal* 26 (11), S. 1115–1140. DOI: 10.1093/eurheartj/ehi204.
- Takenaka, Katsu; Sakamoto, Tsuguya; Amano, Keiko; Oku, Junji; Fujinami, Kenji; Murakami, Tohru et al. (1988): Left ventricular filling determined by Doppler echocardiography in diabetes mellitus. In: *The American Journal of Cardiology* 61 (13), S. 1140–1143. DOI: 10.1016/0002-9149(88)90149-X.
- Talbott, Evelyn O.; Zborowski, Jeanne V.; Rager, Judith R.; Kip, Kevin E.; Xu, Xiaohui; Orchard, Trevor J. (2007): Polycystic ovarian syndrome (PCOS): a significant contributor to the overall burden of type 2 diabetes in women. In: *Journal of women's health* (2002) 16 (2), S. 191–197. DOI: 10.1089/jwh.2006.0098.
- Tan, K. H.; Cossart, L. de; Edwards, P. R. (2000): Exercise training and peripheral vascular disease. In: *British Journal of Surgery* 87 (5), S. 553–562. DOI: 10.1046/j.1365-2168.2000.01445.x.
- Tan, Ricardo A.; Spector, Sheldon L. (2002): Exercise-induced asthma: diagnosis and management. In: *Annals of Allergy, Asthma & Immunology* 89 (3), S. 226–236. DOI: 10.1016/S1081-1206(10)61948-5.
- Tarumi, N.; Iwasaka, T.; Takahashi, N.; Sugiura, T.; Morita, Y.; Sumimoto, T. et al. (1993): Left ventricular diastolic filling properties in diabetic patients during isometric exercise. In: *Cardiology* 83 (5-6), S. 316–323. DOI: 10.1159/000175987.
- TASC (2000): Management of peripheral arterial disease. In: *Eur J Vasc Endovasc Surg* 19, S1.
- Taylor (2014): Exercise-based rehabilitation for heart failure. In: *Cochrane Database Syst Rev* 4, CD003331.
- Taylor, Rod S.; Brown, Allan; Ebrahim, Shah; Jolliffe, Judith; Noorani, Hussein; Rees, Karen et al. (2004): Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. In: *The American Journal of Medicine* 116 (10), S. 682–692. DOI: 10.1016/j.amjmed.2004.01.009.
- Taylor, C. B.; Jatulis, D. E.; Winkleby, M. A.; Rockhill, B. J.; Kraemer, H. C. (1994): Effects of life-style on body mass index change. In: *Epidemiology* 5 (6), S. 599–603. DOI: 10.1097/00001648-199411000-00007.
- Thelle, Dag S.; FØRDE, OLAV H.; Try, Kenneth; Lehmann, Egil H. (1976): The Tromsø Heart Study. In: *Acta Medica Scandinavica* 200 (1-6), S. 107–118. DOI: 10.1111/j.0954-6820.1976.tb08204.x.
- Theorell (2006): Stress and cardiovascular disease.
- Thirlaway, Kathryn; Benton, David (1992): Participation in physical activity and cardiovascular fitness have different effects on mental health and mood. In: *Journal of Psychosomatic Research* 36 (7), S. 657–665. DOI: 10.1016/0022-3999(92)90055-7.
- Thomas (2006): Exercise for type 2 diabetes mellitus. In: *Cochrane Database Syst Rev* 19, CD002968.
- Thomas, E. L.; Brynes, A. E.; McCarthy, J.; Goldstone, A. P.; Hajnal, J. V.; Saeed, N. et al. (2000): Preferential loss of visceral fat following aerobic exercise, measured by magnetic resonance imaging. In: *Lipids* 35 (7), S. 769–776. DOI: 10.1007/s11745-000-0584-0.
- Thompson, Paul D.; Buchner, David; Pina, Ileana L.; Balady, Gary J.; Williams, Mark A.; MARCUS, BESS H. et al. (2003): Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease: a statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity). In: *Circulation* 107 (24), S. 3109–3116. DOI: 10.1161/01.CIR.0000075572.40158.77.
- Thomson, Rebecca L.; Buckley, Jonathan D.; Noakes, Manny; Clifton, Peter M.; Norman, Robert J.; Brinkworth, Grant D. (2008): The effect of a hypocaloric diet with and without exercise training on body composition, cardiometabolic risk profile, and reproductive function in overweight and obese women with polycystic ovary syndrome. In: *The Journal of clinical endocrinology and metabolism* 93 (9), S. 3373–3380. DOI: 10.1210/jc.2008-0751.

- Thomson, R. L.; Buckley, J. D.; Brinkworth, G. D. (2011): Exercise for the treatment and management of overweight women with polycystic ovary syndrome: a review of the literature. In: *Obesity Reviews* 12 (5), e202-10. DOI: 10.1111/j.1467-789X.2010.00758.x.
- Thune, I. (1998): Physical exercise in rehabilitation program for cancer patients? In: *Journal of alternative and complementary medicine (New York, N.Y.)* 4 (2), S. 205–207. DOI: 10.1089/acm.1998.4.205.
- Thune, I.; Furberg, A. S. (2001): Physical activity and cancer risk: dose-response and cancer, all sites and site-specific. In: *Medicine & Science in Sports & Exercise* 33 (6 Suppl), S530-50; discussion S609-10. DOI: 10.1097/00005768-200106001-00025.
- Tipton (1999): Exercise and the heart in health and disease.
- Tjønnå, Arnt Erik; Lee, Sang Jun; Rognum, Øivind; Stølen, Tomas O.; Bye, Anja; Haram, Per Magnus et al. (2008): Aerobic interval training versus continuous moderate exercise as a treatment for the metabolic syndrome: a pilot study. In: *Circulation* 118 (4), S. 346–354. DOI: 10.1161/CIRCULATIONAHA.108.772822.
- Tolmunen, T.; Laukkanen, J. A.; Hintikka, J.; Kurl, S.; Viinamäki, H.; Salonen, R. et al. (2006): Low maximal oxygen uptake is associated with elevated depressive symptoms in middle-aged men. In: *European Journal of Epidemiology* 21 (9), S. 701–706. DOI: 10.1007/s10654-006-9038-5.
- Tomlinson, Deborah; Diorio, Caroline; Beyene, Joseph; Sung, Lillian (2014): Effect of exercise on cancer-related fatigue: a meta-analysis. In: *American Journal of Physical Medicine & Rehabilitation* 93 (8), S. 675–686. DOI: 10.1097/PHM.0000000000000083.
- Torres Lacomba, María; Yuste Sánchez, María José; Zapico Goñi, Alvaro; Prieto Merino, David; Mayoral del Moral, Orlando; Cerezo Téllez, Ester; Minayo Mogollón, Elena (2010): Effectiveness of early physiotherapy to prevent lymphoedema after surgery for breast cancer: randomised, single blinded, clinical trial. In: *BMJ* 340 (jan12 1), b5396. DOI: 10.1136/bmj.b5396.
- Tran (1983): The effects of exercise on blood lipids and lipoproteins: a meta-analysis of studies. In: *Med Sci Sports Exerc* 15, S. 393.
- Tran, Zung Vu (1985): Differential Effects of Exercise on Serum Lipid and Lipoprotein Levels Seen With Changes in Body Weight. In: *JAMA* 254 (7), S. 919. DOI: 10.1001/jama.1985.03360070057023.
- Troosters, T.; Langer, D.; Vrijsen, B.; Segers, J.; Wouters, K.; Janssens, W. et al. (2009): Skeletal muscle weakness, exercise tolerance and physical activity in adults with cystic fibrosis. In: *The European respiratory journal* 33 (1), S. 99–106. DOI: 10.1183/09031936.00091607.
- Trovati, M.; Carta, Q.; Cavalot, F.; Vitali, S.; Banaudi, C.; Lucchina, P. G. et al. (1984): Influence of physical training on blood glucose control, glucose tolerance, insulin secretion, and insulin action in non-insulin-dependent diabetic patients. In: *Diabetes Care* 7 (5), S. 416–420. DOI: 10.2337/diacare.7.5.416.
- Tu, Rong-Hui; Zeng, Zhi-Yu; Zhong, Guo-Qiang; Wu, Wei-Feng; Lu, Yun-Juan; Bo, Zhan-Dong et al. (2014): Effects of exercise training on depression in patients with heart failure: a systematic review and meta-analysis of randomized controlled trials. In: *European Journal of Heart Failure* 16 (7), S. 749–757. DOI: 10.1002/ehf.101.
- Tudor-Locke (2000): Formative evaluation of The First Step Program: a practical intervention to increase daily physical activity. In: *Can J Diabetes Care* 24, S. 34.
- Tudor-Locke, C.; Bell, R. C.; Myers, A. M.; Harris, S. B.; Ecclestone, N. A.; Lauzon, N.; Rodger, N. W. (2004): Controlled outcome evaluation of the First Step Program: a daily physical activity intervention for individuals with type II diabetes. In: *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 28 (1), S. 113–119. DOI: 10.1038/sj.ijo.0802485.
- Tudor-Locke, C. E.; Myers, A. M.; Rodger, N. W. (2001): Development of a theory-based daily activity intervention for individuals with type 2 diabetes. In: *The Diabetes Educator* 27 (1), S. 85–93. DOI: 10.1177/014572170102700110.
- Tudor-Locke, Catrine E.; Myers, Anita M.; Bell, Rhonda C.; Harris, Stewart B.; Wilson Rodger, N. (2002): Preliminary outcome evaluation of the First Step Program: a daily physical activity intervention for individuals with type 2 diabetes. In: *Patient Education and Counseling* 47 (1), S. 23–28. DOI: 10.1016/S0738-3991(01)00169-0.
- Tulder (2000): Exercise therapy for low back pain. In: *Cochrane Database Syst Rev*, CD000335.
- Tuomilehto, J.; Lindström, J.; Eriksson, J. G.; Valle, T. T.; Hämäläinen, H.; Ilanne-Parikka, P. et al. (2001): Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. In: *The New England journal of medicine* 344 (18), S. 1343–1350. DOI: 10.1056/NEJM200105033441801.
- Tuominen, J. A.; Karonen, S. L.; Melamies, L.; Bolli, G.; Koivisto, V. A. (1995): Exercise-induced hypoglycaemia in IDDM patients treated with a short-acting insulin analogue. In: *Diabetologia* 38 (1), S. 106–111. DOI: 10.1007/BF02369359.
- Turner, Sian; Eastwood, Peter; Cook, Angus; Jenkins, Sue (2011): Improvements in symptoms and quality of life following exercise training in older adults with moderate/severe persistent asthma. In: *Respiration* 81 (4), S. 302–310. DOI: 10.1159/000315142.
- U.S. Department of Health and Human Services (1996): Physical activity and health: a report of the surgeon general.
- Umpierre, Daniel; Ribeiro, Paula A. B.; Kramer, Caroline K.; Leitão, Cristiane B.; Zucatti, Alessandra T. N.; Azevedo, Mirela J. et al. (2011): Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. In: *JAMA* 305 (17), S. 1790–1799. DOI: 10.1001/jama.2011.576.



- VALENTI, Marco; PORZIO, Giampiero; AIELLI, Federica; VERNA, Lucilla; CANNITA, Katia; MANNO, Renato et al. (2008): Physical exercise and quality of life in breast cancer survivors. In: *International Journal of Medical Sciences* 5 (1), S. 24–28. DOI: 10.7150/ijms.5.24.
- van de Weert-van Leeuwen, Pauline B.; Hulzebos, Hendrikus J.; Werkman, Maarten S.; Michel, Sabine; Vijftigschild, Lodewijk A. W.; van Meegen, Marit A. et al. (2014): Chronic inflammation and infection associate with a lower exercise training response in cystic fibrosis adolescents. In: *Respiratory Medicine* 108 (3), S. 445–452. DOI: 10.1016/j.rmed.2013.08.012.
- van den Akker, Lizanne Eva; Heine, Martin; van der Veldt, Nikki; Dekker, Joost; Groot, Vincent de; Beckerman, Heleen (2015): Feasibility and Safety of Cardiopulmonary Exercise Testing in Multiple Sclerosis: A Systematic Review. In: *Arch Phys Med Rehabil* 96 (11), S. 2055–2066. DOI: 10.1016/j.apmr.2015.04.021.
- van den Ende, C. H.; Breedveld, F. C.; Le Cessie, S.; Dijkmans, B. A.; Mug, A. W. de; Hazes, J. M. (2000): Effect of intensive exercise on patients with active rheumatoid arthritis: a randomised clinical trial. In: *Annals of the Rheumatic Diseases* 59 (8), S. 615–621. DOI: 10.1136/ard.59.8.615.
- van den Ende, C. H.; Hazes, J. M.; Le Cessie, S.; Mulder, W. J.; Belfor, D. G.; Breedveld, F. C.; Dijkmans, B. A. (1996): Comparison of high and low intensity training in well controlled rheumatoid arthritis. Results of a randomised clinical trial. In: *Annals of the Rheumatic Diseases* 55 (11), S. 798–805. DOI: 10.1136/ard.55.11.798.
- van der Poest Clement, E.; van der Wiel, H.; Patka, P.; Roos, J.C; Lips, P. (1999): Long-term consequences of fracture of the lower leg: cross-sectional study and long-term longitudinal follow-up of bone mineral density in the hip after fracture of lower leg. In: *Bone* 24 (2), S. 131–134. DOI: 10.1016/S8756-3282(98)00153-7.
- van der Wiel, H. E.; Lips, P.; Nauta, J.; Patka, P.; Haarman, H. J.; Teule, G. J. (1994): Loss of bone in the proximal part of the femur following unstable fractures of the leg. In: *The Journal of Bone & Joint Surgery* 76 (2), S. 230–236. DOI: 10.2106/00004623-199402000-00009.
- van Doorn, Nancy (2010): Exercise programs for children with cystic fibrosis: a systematic review of randomized controlled trials. In: *Disability and Rehabilitation* 32 (1), S. 41–49. DOI: 10.3109/09638280902991842.
- van Janse Rensburg, Dina C.; Ker, James A.; Grant, Catharina C.; Fletcher, Lizelle (2012): Effect of exercise on cardiac autonomic function in females with rheumatoid arthritis. In: *Clinical Rheumatology* 31 (8), S. 1155–1162. DOI: 10.1007/s10067-012-1985-5.
- van Middelkoop, Marienke; Rubinstein, Sidney M.; Kuijpers, Ton; Verhagen, Arianne P.; Ostelo, Raymond; Koes, Bart W.; van Tulder, Maurits W. (2011): A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. In: *European Spine Journal* 20 (1), S. 19–39. DOI: 10.1007/s00586-010-1518-3.
- van Tol, Benno A. F.; Huijsmans, Rosalie J.; Kroon, Dineke W.; Schothorst, Maaïke; Kwakkel, Gert (2006): Effects of exercise training on cardiac performance, exercise capacity and quality of life in patients with heart failure: a meta-analysis. In: *European Journal of Heart Failure* 8 (8), S. 841–850. DOI: 10.1016/j.ejheart.2006.02.013.
- Vancampfort, D.; Knapen, J.; Probst, M.; Scheewe, T.; Remans, S.; Hert, M. de (2012): A systematic review of correlates of physical activity in patients with schizophrenia. In: *Acta Psychiatrica Scandinavica* 125 (5), S. 352–362. DOI: 10.1111/j.1600-0447.2011.01814.x.
- Vancampfort, Davy; Probst, Michel; Helvik Skjaerven, Liv; Catalán-Matamoros, Daniel; Lundvik-Gyllensten, Amanda; Gómez-Conesa, Antonia et al. (2012): Systematic review of the benefits of physical therapy within a multidisciplinary care approach for people with schizophrenia. In: *Physical Therapy* 92 (1), S. 11–23. DOI: 10.2522/ptj.20110218.
- Vanninen, E.; Uusitupa, M.; Siitonen, O.; Laitinen, J.; Länsimies, E. (1992): Habitual physical activity, aerobic capacity and metabolic control in patients with newly-diagnosed type 2 (non-insulin-dependent) diabetes mellitus: effect of 1-year diet and exercise intervention. In: *Diabetologia* 35 (4), S. 340–346. DOI: 10.1007/BF00401201.
- Varlotta, L. (1998): Management and care of the newly diagnosed patient with cystic fibrosis. In: *Current Opinion in Pulmonary Medicine* 4 (6), S. 311–318. DOI: 10.1097/00063198-199811000-00001.
- Vergheze, J.; LeValley, A.; Derby, C.; Kuslansky, G.; Katz, M.; Hall, C. et al. (2006): Leisure activities and the risk of amnesic mild cognitive impairment in the elderly. In: *Neurology* 66 (6), S. 821–827. DOI: 10.1212/01.wnl.0000202520.68987.48.
- Vergheze, Joe; Lipton, Richard B.; Katz, Mindy J.; Hall, Charles B.; Derby, Carol A.; Kuslansky, Gail et al. (2003): Leisure activities and the risk of dementia in the elderly. In: *New England Journal of Medicine* 348 (25), S. 2508–2516. DOI: 10.1056/NEJMoa022252.
- Veves, A.; Saouaf, R.; Donaghue, V. M.; Mullooly, C. A.; Kistler, J. A.; Giurini, J. M. et al. (1997): Aerobic exercise capacity remains normal despite impaired endothelial function in the micro- and macrocirculation of physically active IDDM patients. In: *Diabetes* 46 (11), S. 1846–1852. DOI: 10.2337/diab.46.11.1846.
- Vibek (1991): Respiratory care.
- Vries, Sybolt O. de; Visser, Karen; Vries, Jelle A. de; Wong, John B.; Donaldson, Magruder C.; Hunink, M. G. Myriam (2002): Intermittent claudication: cost-effectiveness of revascularization versus exercise therapy. In: *Radiology* 222 (1), S. 25–36. DOI: 10.1148/radiol.2221001743.

- Vroomen, P. C.; Krom, M. C. de; Wilmlink, J. T.; Kester, A. D.; Knottnerus, J. A. (1999): Lack of effectiveness of bed rest for sciatica. In: *The New England Journal of Medicine* 340 (6), S. 418–423. DOI: 10.1056/NEJM199902113400602.
- Wadden, Thomas A.; Vogt, Renee A.; Foster, Gary D.; Anderson, Drew A. (1998): Exercise and the maintenance of weight loss: 1-year follow-up of a controlled clinical trial. In: *Journal of Consulting and Clinical Psychology* 66 (2), S. 429–433. DOI: 10.1037/0022-006X.66.2.429.
- Wadell, K.; Henriksson-Larsén, K.; Lundgren, R. (2001): Physical training with and without oxygen in patients with chronic obstructive pulmonary disease and exercise-induced hypoxaemia. In: *Journal of Rehabilitation Medicine* 33 (5), S. 200–205. DOI: 10.1080/165019701750419581.
- Walker, K. Z.; Piers, L. S.; Putt, R. S.; Jones, J. A.; O'Dea, K. (1999): Effects of regular walking on cardiovascular risk factors and body composition in normoglycemic women and women with type 2 diabetes. In: *Diabetes Care* 22 (4), S. 555–561. DOI: 10.2337/diacare.22.4.555.
- Wallace, B. A.; Cumming, R. G. (2000): Systematic review of randomized trials of the effect of exercise on bone mass in pre- and postmenopausal women. In: *Calcified Tissue International* 67 (1), S. 10–18. DOI: 10.1007/s00223001089.
- Wallberg-Henriksson, H.; Gunnarsson, R.; Henriksson, J.; Ostman, J.; Wahren, J. (1984): Influence of physical training on formation of muscle capillaries in type I diabetes. In: *Diabetes* 33 (9), S. 851–857. DOI: 10.2337/diab.33.9.851.
- Wallberg-Henriksson, H.; Gunnarsson, R.; Rössner, S.; Wahren, J. (1986): Long-term physical training in female type 1 (insulin-dependent) diabetic patients: absence of significant effect on glycaemic control and lipoprotein levels. In: *Diabetologia* 29 (1), S. 53–57. DOI: 10.1007/BF02427281.
- Waller, Benjamin; Ogonowska-Slodownik, Anna; Vitor, Manuel; Lambeck, Johan; Daly, Daniel; Kujala, Urho M.; Heinonen, Ari (2014): Effect of therapeutic aquatic exercise on symptoms and function associated with lower limb osteoarthritis: systematic review with meta-analysis. In: *Physical Therapy* 94 (10), S. 1383–1395. DOI: 10.2522/ptj.20130417.
- Walsh (1995): A 54-month evaluation of a popular very low calorie diet program. In: *J Fam Pract* 41, S. 231.
- Wang, Chong-Wen; Chan, Celia H. Y.; Ho, Rainbow T. H.; Chan, Jessie S. M.; Ng, Siu-Man; Chan, Cecilia L. W. (2014): Managing stress and anxiety through qigong exercise in healthy adults: a systematic review and meta-analysis of randomized controlled trials. In: *BMC Complementary and Alternative Medicine* 14 (1), S. 8. DOI: 10.1186/1472-6882-14-8.
- Wang, Xue-Qiang; Zheng, Jie-Jiao; Yu, Zhuo-Wei; Bi, Xia; Lou, Shu-Jie; Liu, Jing et al. (2012): A meta-analysis of core stability exercise versus general exercise for chronic low back pain. In: *PLoS ONE* 7 (12), e52082. DOI: 10.1371/journal.pone.0052082.
- Wannamethee, G.; Shaper, A. G. (1992): Physical activity and stroke in British middle aged men. In: *BMJ* 304 (6827), S. 597–601. DOI: 10.1136/bmj.304.6827.597.
- Wasserman, D. H.; Zinman, B. (1994): Exercise in individuals with IDDM. In: *Diabetes Care* 17 (8), S. 924–937. DOI: 10.2337/diacare.17.8.924.
- Wegner, Mirko; Helmich, Ingo; Machado, Sergio; Nardi, Antonio E.; Arias-Carrion, Oscar; Budde, Henning (2014): Effects of exercise on anxiety and depression disorders: review of meta-analyses and neurobiological mechanisms. In: *CNS & neurological disorders drug targets* 13 (6), S. 1002–1014. DOI: 10.2174/1871527313666140612102841.
- Wei, M.; Gibbons, L. W.; Kampert, J. B.; Nichaman, M. Z.; Blair, S. N. (2000): Low cardiorespiratory fitness and physical inactivity as predictors of mortality in men with type 2 diabetes. In: *Annals of Internal Medicine* 132 (8), S. 605–611. DOI: 10.7326/0003-4819-132-8-200004180-00002.
- Wendel-Vos, G. C. W.; Schuit, A. J.; Feskens, E. J. M.; Boshuizen, H. C.; Verschuren, W. M. M.; Saris, W. H. M.; Kromhout, D. (2004): Physical activity and stroke. A meta-analysis of observational data. In: *International Journal of Epidemiology* 33 (4), S. 787–798. DOI: 10.1093/ije/dyh168.
- Westby (2000): A randomized controlled trial to evaluate the effectiveness of an exercise program in women with rheumatoid arthritis taking low dose prednisone. In: *J Rheumatol* 27, S. 1674.
- Whelton, Seamus P.; Chin, Ashley; Xin, Xue; He, Jiang (2002): Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. In: *Annals of Internal Medicine* 136 (7), S. 493–503. DOI: 10.7326/0003-4819-136-7-200204020-00006.
- Wickham, C. A.; Walsh, K.; Cooper, C.; Barker, D. J.; Margetts, B. M.; Morris, J.; Bruce, S. A. (1989): Dietary calcium, physical activity, and risk of hip fracture: a prospective study. In: *BMJ* 299 (6704), S. 889–892. DOI: 10.1136/bmj.299.6704.889.
- Wiesel, S. W.; Cuckler, J. M.; Deluca, F.; Jones, F.; Zeide, M. S.; Rothman, R. H. (1980): Acute low-back pain. An objective analysis of conservative therapy. In: *Spine* 5 (4), S. 324–330. DOI: 10.1097/00007632-198007000-00006.
- Wiesinger (2001): Health related quality of life in patients with long-standing insulin dependent (type 1) diabetes mellitus: benefits of regular physical training. In: *Wien Klin Wochenschr* 113, S. 670.
- Wild, Sarah; Roglic, Gojka; Green, Anders; Sicree, Richard; King, Hilary (2004): Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. In: *Diabetes Care* 27 (5), S. 1047–1053. DOI: 10.2337/diacare.27.5.1047.

- Wilkes, Donna L.; Schneiderman, Jane E.; Nguyen, Thanh; Heale, Liane; Moola, Fiona; Ratjen, Felix et al. (2009): Exercise and physical activity in children with cystic fibrosis. In: *Paediatric Respiratory Reviews* 10 (3), S. 105–109. DOI: 10.1016/j.prrv.2009.04.001.
- Wilkinson (1995): Does 48 hours' bed rest influence the outcome of acute low back pain. In: *Br J Gen Pract* 45, S. 481.
- Williams (2010): Preventing Alzheimer's disease and cognitive decline. In: *Evid Rep Technol Assess (Full Rep)*.
- Williamson (1993): Recreational physical activity and ten-year weight change in a US national cohort. In: *Int J Obes Relat Metab Disord* 17, S. 279.
- Wilson, J. R.; Mancini, D. M.; Dunkman, W. B. (1993): Exertional fatigue due to skeletal muscle dysfunction in patients with heart failure. In: *Circulation* 87 (2), S. 470–475. DOI: 10.1161/01.CIR.87.2.470.
- WILSON, MARCUS G.; MICHET, CLEMENT J.; ILSTRUP, DUANE M.; MELTON III, L. JOSEPH (1990): Idiopathic Symptomatic Osteoarthritis of the Hip and Knee: A Population-Based Incidence Study. In: *Mayo Clinic Proceedings* 65 (9), S. 1214–1221. DOI: 10.1016/S0025-6196(12)62745-1.
- Wind, J.; Koelemay, M. J. W. (2007): Exercise therapy and the additional effect of supervision on exercise therapy in patients with intermittent claudication. Systematic review of randomised controlled trials. In: *European Journal of Vascular and Endovascular Surgery* 34 (1), S. 1–9. DOI: 10.1016/j.ejvs.2006.12.030.
- Wing, Rena R.; Bolin, Paula; Brancati, Frederick L.; Bray, George A.; Clark, Jeanne M.; Coday, Mace et al. (2013): Cardiovascular effects of intensive lifestyle intervention in type 2 diabetes. In: *New England Journal of Medicine* 369 (2), S. 145–154. DOI: 10.1056/NEJMoA1212914.
- Wing, R. R.; Epstein, L. H.; Paternostro-Bayles, M.; Kriska, A.; Nowalk, M. P.; Gooding, W. (1988): Exercise in a behavioural weight control programme for obese patients with Type 2 (non-insulin-dependent) diabetes. In: *Diabetologia* 31 (12), S. 902–909. DOI: 10.1007/BF00265375.
- Winocour, Peter H.; Durrington, Paul N.; Bhatnagar, Deepak; Mbewu, Anthony D.; Ishola, Monica; Mackness, Michael; Arrol, Sharon (1992): A cross-sectional evaluation of cardiovascular risk factors in coronary heart disease associated with Type 1 (insulin-dependent) diabetes mellitus. In: *Diabetes Research and Clinical Practice* 18 (3), S. 173–184. DOI: 10.1016/0168-8227(92)90143-F.
- Wisløff, Ulrik; Støylen, Asbjørn; Loennechen, Jan P.; Bruvold, Morten; Rognum, Øivind; Haram, Per Magnus et al. (2007): Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. In: *Circulation* 115 (24), S. 3086–3094. DOI: 10.1161/CIRCULATIONAHA.106.675041.
- Wolfe, F.; Mitchell, D. M.; Sibley, J. T.; Fries, J. F.; Bloch, D. A.; Williams, C. A. et al. (1994): The mortality of rheumatoid arthritis. In: *Arthritis and rheumatism* 37 (4), S. 481–494. DOI: 10.1002/art.1780370408.
- Wolin, K. Y.; Yan, Y.; Colditz, G. A.; Lee, I-M (2009): Physical activity and colon cancer prevention: a meta-analysis. In: *British Journal of Cancer* 100 (4), S. 611–616. DOI: 10.1038/sj.bjc.6604917.
- Wu, T.; Gao, X.; Chen, M.; van Dam, R. M. (2009): Long-term effectiveness of diet-plus-exercise interventions vs. diet-only interventions for weight loss: a meta-analysis. In: *Obesity Reviews* 10 (3), S. 313–323. DOI: 10.1111/j.1467-789X.2008.00547.x.
- Wu, Mei-Kuei; Wang, Chin-Kun; Bai, Ya-Mei; Huang, Chih-Yang; Lee, Shin-Da (2007): Outcomes of obese, clozapine-treated inpatients with schizophrenia placed on a six-month diet and physical activity program. In: *Psychiatric services (Washington, D.C.)* 58 (4), S. 544–550. DOI: 10.1176/ps.2007.58.4.544.
- Wu, Ren-Rong; Zhao, Jing-Ping; Jin, Hua; Shao, Ping; Fang, Mao-Sheng; Guo, Xiao-Feng et al. (2008): Lifestyle intervention and metformin for treatment of antipsychotic-induced weight gain: a randomized controlled trial. In: *JAMA* 299 (2), S. 185–193. DOI: 10.1001/jama.2007.56-b.
- Yaffe, K.; Barnes, D.; Nevitt, M.; Lui, L. Y.; Covinsky, K. (2001): A prospective study of physical activity and cognitive decline in elderly women: women who walk. In: *Archives of Internal Medicine* 161 (14), S. 1703–1708. DOI: 10.1001/archinte.161.14.1703.
- Yaffe, K.; Fiocco, A. J.; Lindquist, K.; Vittinghoff, E.; Simonsick, E. M.; Newman, A. B. et al. (2009): Predictors of maintaining cognitive function in older adults: the Health ABC study. In: *Neurology* 72 (23), S. 2029–2035. DOI: 10.1212/WNL.0b013e3181a92c36.
- Yamanouchi, K.; Shinozaki, T.; Chikada, K.; Nishikawa, T.; Ito, K.; Shimizu, S. et al. (1995): Daily walking combined with diet therapy is a useful means for obese NIDDM patients not only to reduce body weight but also to improve insulin sensitivity. In: *Diabetes Care* 18 (6), S. 775–778. DOI: 10.2337/diacare.18.6.775.
- Yang, Yea-Ru; Lee, Ya-Yun; Cheng, Shih-Jung; Wang, Ray-Yau (2010): Downhill walking training in individuals with Parkinson's disease: a randomized controlled trial. In: *American Journal of Physical Medicine & Rehabilitation* 89 (9), S. 706–714. DOI: 10.1097/PHM.0b013e3181e721c5.
- Yang, Zuyao; Scott, Catherine A.; Mao, Chen; Tang, Jinling; Farmer, Andrew J. (2014): Resistance exercise versus aerobic exercise for type 2 diabetes: a systematic review and meta-analysis. In: *Sports medicine (Auckland, N.Z.)* 44 (4), S. 487–499. DOI: 10.1007/s40279-013-0128-8.

Yasuda (1992): Systolic and diastolic left ventricular dysfunction in middle-aged asymptomatic non-insulin-dependent diabetics. In: *J Cardiol* 22, S. 427.

Yii, Ming Fong; Lim, Chi Eung Danforn; Luo, Xiping; Wong, Wu Shun Felix; Cheng, Nga Chong Lisa; Zhan, Xinlin (2009): Polycystic ovarian syndrome in adolescence. In: *Gynecological Endocrinology* 25 (10), S. 634–639. DOI: 10.1080/09513590903015551.

Yki-Järvinen, H.; DeFronzo, R. A.; Koivisto, V. A. (1984): Normalization of insulin sensitivity in type I diabetic subjects by physical training during insulin pump therapy. In: *Diabetes Care* 7 (6), S. 520–527. DOI: 10.2337/diacare.7.6.520.

Yoshitake, T.; Kiyohara, Y.; Kato, I.; Ohmura, T.; Iwamoto, H.; Nakayama, K. et al. (1995): Incidence and risk factors of vascular dementia and Alzheimer's disease in a defined elderly Japanese population: the Hisayama Study. In: *Neurology* 45 (6), S. 1161–1168. DOI: 10.1212/WNL.45.6.1161.

Zanuso, Silvano; Jimenez, A.; Pugliese, G.; Corigliano, G.; Balducci, S. (2010): Exercise for the management of type 2 diabetes: a review of the evidence. In: *Acta Diabetologica* 47 (1), S. 15–22. DOI: 10.1007/s00592-009-0126-3.

Zavaroni, I.; Bonini, L.; Gasparini, P.; Barilli, A. L.; Zuccarelli, A.; Dall'Aglio, E. et al. (1999): Hyperinsulinemia in a normal population as a predictor of non—insulin-dependent diabetes mellitus, hypertension, and coronary heart disease: The barilla factory revisited. In: *Metabolism* 48 (8), S. 989–994. DOI: 10.1016/S0026-0495(99)90195-6.

Zhang, W.; Nuki, G.; Moskowitz, R. W.; Abramson, S.; Altman, R. D.; Arden, N. K. et al. (2010): OARSI recommendations for the management of hip and knee osteoarthritis: part III: Changes in evidence following systematic cumulative update of research published through January 2009. In: *Osteoarthritis and Cartilage* 18 (4), S. 476–499. DOI: 10.1016/j.joca.2010.01.013.